



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



6044 Cornerstone Court W, Ste E
San Diego, CA 92121
Tel: 1.858.829.3082
Fax: 1.858.481.8694
Email: info@bpsbioscience.com

BCA1 (CXCL13)

Catalog #: 90104-B (20 µg)
Lot #: 150723

Description: Recombinant human BCA1 (B cell- attracting chemokine 1), also known as CXCL13 or BLC, is a disulfide-linked monomeric protein consisting of 88 amino acids and migrates as an approximately 10 kDa protein under reducing and non-reducing conditions. Human CXCL13, generated by the proteolytic removal of the signal peptide and propeptide, has a calculated molecular mass of ~10 kDa.

Source: Optimized DNA sequence encoding human BCA1 mature domain was expressed in *E. coli*.

Formulation: Lyophilized from a 0.2 µm filtered solution containing 20 mM phosphate buffer, pH 7.5, 100 mM NaCl.

Reconstitution: Reconstitute at 0.1-1.0 mg/mL in distilled water. This solution can then be diluted into other buffers. To maximize product collection from vial surface, vortex briefly and then spin down to recollect the liquid.

Storage: The lyophilized protein is stable for at least 2 years from date of receipt when stored at -20°C. Upon reconstitution, store in working aliquots at 2 - 8° C for up to one month, or at -20°C for up to six months, in the presence of a carrier protein. Avoid repeated freeze/thaw cycles.

Purity: >98%, as determined by SDS-PAGE and HPLC.

Endotoxin Level: Endotoxin level was found to be < 0.1 ng/µg (1EU/µg), using the LAL gel clot method.

Biological Activity: Determined by its ability to chemoattract human CXCR5 transfected mouse BaF3 B cells with ED50 >20 ng/ml.

Protein Sequence:

VLEVYYTSLRCRCVQESSVFIPRRFID
RIQILPRGNGCPRKEIIVWKKNKSIKCV
DPQAEWIQRMMEVLRKRSSSTLPVPV
FKRKIP

References:

1. *Neurology*, Dec 2009; **73**: 2003 - 2010.
2. *Nephrol. Dial. Transplant.*, Dec 2009; **24**: 3708 - 3712.