

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in





Revised: January 21, 2019

Product Information

Rabbit Anti-Goat IgG (H+L), Whole Antibody, CF® Dye Conjugates

CF® Dye	Ex/Em	Catalog No.	Size
CF®488A	490/515 nm	20021-1	50 uL
		20021	500 uL
		20021-1mg	1 mg (lyophilized)
CF®543	541/560 nm	20315-1	50 uL
		20315	500 uL
		20315-1mg	1 mg (lyophilized)
CF®555	555/565 nm	20031-1	50 uL
		20031	500 uL
		20031-1mg	1 mg (lyophilized)
CF®568	562/583 nm	20107-1	50 uL
		20107	500 uL
		20107-1mg	1 mg (lyophilized)
CF®594	593/614 nm	20117-1	50 uL
		20117	500 uL
		20117-1mg	1 mg (lyophilized)
CF®633	630/650 nm	20128-1	50 uL
		20128	500 uL
		20128-1mg	1 mg (lyophilized)
CF®640R	642/662 nm	20090-1	50 uL
		20090	500 uL
		20090-1mg	1 mg (lyophilized)
CF®647	650/665 nm	20049-1	50 uL
		20049	500 uL
		20049-1mg	1 mg (lyophilized)
CF®680	681/698 nm	20068-1	50 uL
		20068	500 uL
		20068-1mg	1 mg (lyophilized)

Concentration

Liquid format: 2 mg/mL in pH \sim 7.4 PBS containing 50% glycerol, 2 mg/ml bovine serum albumin (IgG-free and protease-free) and 0.05% sodium azide.

Lyophilized format (after reconstitution): 2 mg/mL in pH ~7.4 PBS containing 15 mg/mL bovine serum albumin (lgG-free and protease-free) and 20 mg/mL trehalose.

Storage and Handling

Store at -20°C, protected from light. Product is stable for at least 6 months from date of receipt when stored as recommended. Liquid format antibodies contain 50% glycerol and will not freeze at -20°C.

Reconstitution (lyophilized format only): add 0.5 mL dH $_2$ O and mix gently to dissolve. Store at -20°C, protected from light. Aliquot to avoid freeze-thaw cycles. Alternatively, glycerol can be added to the antibody so that it will not freeze at -20°C: add 0.25 mL dH $_2$ O to the lyophilized antibody and mix gently to dissolve, then add 0.25 mL glycerol and mix well. Optional: a preservative may be added, such as 0.05% (final concentration) sodium azide.

Note: storage of the antibody for more than a day at final working dilution is not recommended.

Product Description

Rabbit anti-goat IgG (H+L), whole antibody is an affinity purified rabbit antibody that recognizes goat IgG. This antibody is not highly-cross adsorbed against other species. Visit www.biotium.com to see our highly cross-adsorbed secondary antibodies for multiple labeling.

CF® dyes are superior to Alexa Fluor®, DyLight®, and Cy® dyes for antibody labeling by having combined advantages in brightness, photostability, specificity and novel features ideal for in vivo imaging. For more information about CF® dyes, download our CF® Dye Selection Guide at www.biotium.com.

Recommended Dilution Range

Fluorescence microscopy: 1-2 ug/mL Flow cytometry: 1 ug/10⁶ cells

Near-infrared western detection: 50-100 ng/mL

These concentrations are provided as a starting point for optimization; appropriate dilutions should be determined empirically. Generally IgG conjugates are used in the range of 1-10 ug/mL.

For more detailed protocols for immunofluorescence staining for fluorescence microscopy, flow cytometry, or fluorescence-based western detection, please visit our website.

Related Products

Cat. No.	Product Name	Features	
23001	EverBrite™ Mounting Medium	Excellent protection from photobleaching	
23002	EverBrite™ Mounting Medium with DAPI		
23003	EverBrite™ Hardset Mounting Medium	Compatible with a wide variety of dyes including cyanine dyes Available in wet-set or hardset formulations	
23004	EverBrite™ Hardset Mounting Medium with DAPI	With or without DAPI	
23005	CoverGrip™ Coverslip Sealant	Replaces nail polish for coverslip sealing Won't mix with aqueous mounting media	
23007	TrueBlack™ Lipofuscin Autofluorescence Quencher	Eliminates lipofuscin autofluorescence with less background than Sudan Black B Reduces background from other sources Can be used before or after IF staining	
40061	RedDot™2 Far Red Nuclear Counterstain	Far-red nuclear dye for the Cy®5 channel More specific than Draq®7	
22005	Mini Super ^{HT} Pap Pen 2.5 mm tip, ~400 uses	Create hydrophobic barriers around tissue sections Heat-stable to 120°C Insoluble in aqueous buffers, detergents, alcohol and acetone; can be removed with xylene	
22006	Super HT Pap Pen 4 mm tip, ~800 uses		
22003	Mini-Cell Scrapers	Collect cells from 96-well to 24-well plates Polypropylene, sterile, disposable	
23006	Flow Cytometry Fixation/Permeabilization Kit	Ready-to-use fix/perm buffers for intracellular staining	
22015	Fixation Buffer		
22016	Permeabilization Buffer		
22017	Permeabilization and Blocking Buffer		
22010	10X Fish Gelatin Blocking Agent	Convenient buffers and blocking agents for immunofluorescence or western	
22011	Fish Gelatin Powder		
22014	30% Bovine Serum Albumin Solution		
22002	Tween®-20		

Please visit www.biotium.com to view our full selection of products featuring bright and photostable fluorescent CF® dyes, including Mix-n-Stain™ antibody labeling kits, primary antibody conjugates, streptavidin, phalloidin, and other bioconjugates, as well as conjugates of biotin, HRP, AP, R-PE, APC and PerCP.

Materials from Biotium are sold for research use only, and are not intended for food, drug, household, or cosmetic use.

CF dye technology is covered by pending US and international patents. Alexa Fluor is a registered trademark of Molecular Probes; CYDYE is a registered trademark of GE Healthcare; Draq7 is a registered trademark of Biostatus Ltd; DyLight is a registered trademark of Pierce Biotechnology; TWEEN is a registered trademark of Uniquema Americas LLC.