



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Product Information

CF™ Dye Conjugated Goat Anti-Human IgM (mu chain) Antibodies

Catalog No.	Product Description
20346	CF647 Goat anti-human IgM (mu chain)
20347	CF488A Goat anti-human IgM (mu chain)
20348	CF594 Goat anti-human IgM (mu chain)
20349	CF640R Goat anti-human IgM (mu chain)

Unit size: 50 uL or 250 uL (liquid format)

Concentration: 2 mg/mL in pH ~7.4 PBS containing 50% glycerol, 2 mg/ml bovine serum albumin (IgG-free and protease-free) and 0.05% sodium azide.

Spectral Properties

$\lambda_{abs}/\lambda_{em}$ maxima in pH 7.4 PBS buffer

Conjugate	Abs (nm)	Em (nm)
CF488A	490	515
CF594	593	614
CF640R	642	662
CF647	650	665

Storage and Handling

Store at -20°C, protected from light. Product is stable for at least 6 months from date of receipt when stored as recommended. Liquid format antibodies contain 50% glycerol and will not freeze at -20°C.

Note: storage of the antibody for more than a day at final working dilution is not recommended.

Product Description

Goat anti-human IgM (μ chain) antibody, CF™ dye conjugate, is prepared from antibodies that have been affinity purified and the CF™ labeled conjugate should be suitable for immunofluorescence and other standard immunoassay applications. The antibodies react specifically with IgM heavy chains (μ chains) and not with immunoglobulin light chains. To minimize crossreactivity, it has been adsorbed human IgG prior to conjugation.

General Protocols for Using CF™-Dye Labeled Secondary Antibodies

Recommended Dilution Range

1-10 μ g/mL of the IgM conjugate for most applications (appropriate dilutions of the conjugate should be determined empirically). See other side for example staining protocols.

Immunofluorescence Protocol for Microscopy

There are many methods for immunofluorescence staining. The protocol below is a general guideline for staining cells and should be optimized or modified to obtain the best results for each particular application.

1. Coverslip preparation for adherent cells

- Culture cells on slide chambers or sterile glass coverslips (with poly-L-lysine coating if cells do not adhere well, see below). We recommend 18 x 18 mm square coverslips in 6-well plates or 4-well chamber slides.
- Allow cells to adhere and treat as desired.
- Rinse cells gently with PBS.

2. Coverslip preparation for non-adherent cells

- Coat coverslips with 0.01% poly-L-lysine solution for 10 minutes at room temperature.
- Aspirate the poly-L-lysine solution and allow coverslips to dry completely.
- Centrifuge cells in medium and resuspend in PBS. Transfer cells to coverslips.
- Incubate for 30-60 minutes. Check for adherence by microscope.

3. Fixation and Staining

- Fix with 4% paraformaldehyde/PBS, 15 min.
- Rinse twice with PBS to remove traces of fixative.
- Permeabilize with 0.1 - 0.5% TritonX-100/PBS, 5-10 min.
- Block with blocking agent such as with 5% BSA or normal goat serum in PBS, 30 min.
- Dilute primary antibody in dilution buffer as recommended in the specific product's datasheet. Overlay enough diluted antibody to cover cells on coverslip (150-200 μ L is usually sufficient to cover the surface area) or add to each chamber of the chamber slides. Keep slips covered or in a humidified chamber to avoid evaporation.
- Rinse three times with PBS, 5 min each wash.
- Dilute fluorescent secondary antibody in dilution buffer and incubate for 1 hour at room temperature. General range for IgM conjugates is between 1-10 μ g/mL for most applications. Cell samples without primary antibody incubation is recommended for background control. Keep slips covered or in a humidified chamber to avoid evaporation.
- Rinse three times with PBS, 5 min each wash.
- Additional staining with fluorescent nuclear stains or phalloidins can be done at this step.
- Invert each coverslip onto a pre-cleaned slide with fluorescence anti-fade mounting media. Seal edges with clear polish if desired.
- Store slides in the dark at 4°C.

Staining Protocol for Flow Cytometry

There are many alternative procedures that can be used for specific staining experiments. The protocol below is a general guideline for flow cytometry and should be optimized or modified for each application.

- Aliquot 1 X 10⁶ cells into 12 X 75 mm polypropylene tubes for flow cytometry.
- For intracellular staining, cells can be fixed first to ensure stability of soluble antigens or antigens with short half-lives. We recommend a fix and perm kit from reliable manufacturers. Follow manufacturer's instructions.
- Add the primary antibody or isotype control at the appropriate dilution to the assay tubes. Incubate according to manufacturer's instructions.
- Rinse cells twice by centrifugation with 2-3 mL incubation buffer.
- Decant supernatant and re-suspend the pellet in remaining volume of wash.
- Add fluorescent secondary antibody and incubate for 20-30 minutes. General range for secondary antibodies is between 1-10 μ g/mL for IgM conjugates for most applications.
- Rinse cells twice by centrifugation with 2-3 mL incubation buffer. Centrifuge to collect cells after each wash. Decant supernatant.
- Resuspend cells in 0.5 mL of diluent of choice to analyze on flow cytometer. Acquire data using the correct channel.

Tips and Hints

- No signal or weak fluorescence intensity may suggest the following: (a) insufficient antibody is present for detection, (b) intracellular target was not accessible, (c) excitation sources are not aligned, (d) target protein is not present or expressed at low levels, (e) fluorochrome has faded, and/or (f) primary and secondary antibodies are not compatible.
- High fluorescence intensity may suggest the following: (a) antibody concentration is too high, (b) excess antibody was not washed away efficiently, and/or (c) blocking was inadequate. Increase antibody dilution and washes.

CF™-labeled antibodies can also be used for staining histological sections from paraffin-embedded or frozen tissues.

References

1. Donaldson, J.G. Immunofluorescence staining. (2001) Curr Protoc Cell Biol. Chapter 4: Unit 4.3.
2. Blose, S.H. and Feramisco, J.R. (1983) Fluorescent methods in the analysis of cell structure. Cold Spring Harbour Laboratory.

Useful websites:

www.chroma.com

Related Products

Cat.#	Product Name	Unit Size
40061-T	RedDot™2 Far Red Nuclear Counterstain, 200X in DMSO, Trial Size (15-20 tests)	25 uL
23001	EverBrite™ Mounting Medium	10 mL
23002	EverBrite™ Mounting Medium with DAPI	10 mL
23003	EverBrite™ Hardset Mounting Medium	10 mL
23004	EverBrite™ Hardset Mounting Medium with DAPI	10 mL
23005	CoverGrip™ Coverslip Sealant	15 mL
22005	Mini Super ^{HT} Pap Pen 2.5 mm tip, ~400 uses	1 pen
22006	Super ^{HT} Pap Pen 4 mm tip, ~800 uses	1 pen
22015	Fixation Buffer	100 mL
22016	Permeabilization Buffer	100 mL
22017	Permeabilization and Blocking Buffer	100 mL
22010	10% Fish Gelatin Blocking Buffer	100 mL
22011	Fish Gelatin Powder	2 x 50 g
22014	30% Bovine Serum Albumin Solution	100 mL
22002	Tween®-20	50 mL

Please visit www.biotium.com to view our full selection of products featuring bright and photostable fluorescent CF™ dyes, including streptavidin and other bioconjugates, secondary antibodies, Mix-n-Stain™ antibody labeling kits, and R-PE conjugates. Biotium also offers a variety of apoptosis and cell viability assays for flow cytometry analysis, including mitochondrial membrane potential dyes, fluorescent Annexin V conjugates, and NucView™488 Caspase-3 Substrate for live cells.

CF dye technology is covered by pending U.S. and international patents. Materials from Biotium are sold for research use only, and are not intended for food, drug, household, or cosmetic use.

Tween is a registered trademark of Uniqema Americas LLC.