

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



1,2-Didecanoyl PC

Catalog No. 10009879

CAS Registry No.: 3436-44-0

Formal Name: 1,2-bis(O-decanoyl)-sn-glyceryl-3-

phosphorylcholine

Synonyms: 1,2-Didecanoyl Phosphatidylcholine,


Didecanoyl Lecithin, DPC

 $C_{28}H_{56}NO_8P$ MF: 565.7 FW: **Purity:** ≥98%

Supplied as: A solution in ethanol

Storage: -20°C Stability: ≥2 years

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.



Laboratory Procedures

1,2-Didecanoyl PC (DPC) is supplied as a solution in ethanol. To change the solvent, simply evaporate the ethanol under a gentle stream of nitrogen and immediately add the solvent of choice. Solvents such as ethanol, DMSO, and dimethyl formamide (DMF) purged with an inert gas can be used. The solubility of DPC in ethanol is approximately 30 mg/ml and approximately 20 mg/ml in DMSO and DMF.

DPC is sparingly soluble in aqueous buffers. For maximum solubility in aqueous buffers, the ethanolic solution of DPC should be diluted with the aqueous buffer of choice. DPC has a solubility of approximately 0.5 mg/ml in a 1:1 solution of ethanol:PBS (pH 7.2) using this method. We do not recommend storing the aqueous solution for more than one day.

Description

Phosphatidylcholine species are a common class of phospholipids that comprise the mammalian cell membrane. DPC is a synthetic, less hydrophobic phospholipid that has been found to be useful for enhancing the absorption of peptide drugs and hormones such as insulin. Thus, the bioavailability of intranasally applied human growth hormone is enhanced when coadministered with the absorption enhancer DPC.1

Reference

1. Agerholm, C., Bastholm, L., Johansen, P.B., et al. Epithelial transport and bioavailability of intranasally administered human growth hormone formulated with the absorption enhancers didecanoyl-L-αphosphatidylcholine and α-cyclodextrin in rabbits. J. Pharm. Sci. 83(12), 1706-1711 (1994).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website

Copyright Cayman Chemical Company, 11/30/2021

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM