

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



Vialinin A

Item No. 10010519

CAS Registry No.: 858134-23-3 Formal Name: benzeneacetic acid,

1,1'-(4,4'',5',6'-tetrahydroxy[1,1':4',1''-


terphenyl]-2',3'-diyl) ester

Synonym: Terrestrin A MF: $C_{34}H_{26}O_{8}$ FW: 562.6 **Purity:** ≥95%

 λ_{max} : 263 nm UV/Vis.: Supplied as: A crystalline solid

-20°C Storage: Stability: ≥2 years Item Origin: Synthetic

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.



Laboratory Procedures

Vialinin A is supplied crystalline solid. A stock solution may be made by dissolving the vialinin A in an organic solvent purged with an inert gas. Vialinin A is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide. The solubility of vialinin A in ethanol is approximately 10 mg/ml and approximately 20 mg/ml in DMSO and DMF.

Vialinin A is sparingly soluble in aqueous buffers. For maximum solubility in aqueous buffers, vialinin A should first be dissolved in DMSO and then diluted with the aqueous buffer of choice. Vialinin A has a solubility of approximately 0.25 mg/ml in a 1:3 solution of DMSO:PBS (pH 7.2) using this method. We do not recommend storing the aqueous solution for more than one day.

Description

Vialinin A is a terphenyl compound originally isolated from the fungi T. terrestris and T. vialis. 1,2 Terphenyls, in general, are recognized as strong antioxidants.3 Vialinin A potently inhibits the release of TNF- α (IC₅₀ = 0.09 nM) and IL-4 (IC₅₀ = 2.8 nM), as well as β -hexosaminidase and CCL2 (MCP-1) from IgE-stimulated RBL-2H3 mast cells. 4 Vialinin A does not significantly increase lactate dehydrogenase release from RBL-2H3 mast cells, suggesting low cytotoxicity.⁴

References

- 1. Radulovic, N., Quang, D.N., Hashimoto, T., et al. Terrestrins A-G: p-teraphenyl derivatives from the inedible mushroom Thelephora terrestris. Phytochemistry 66(9), 1052-1059 (2005).
- Xie, C., Koshino, H., Esumi, Y., et al. Vialinin A, a novel 2,2-diphenyl-1-picrylhydrazyl (DPPH) radical scavenger from an edible mushroom in China. Biosci. Biotech. Biochem. 69(12), 2326-2332 (2005).
- 3. Liu, J.-K. Natural terphenyls: Developments since 1877. Chem. Rev. 106(6), 2209-2223 (2006).
- 4. Onose, J., Xie, C., Ye, Y.Q., et al. Vialinin A, a novel potent inhibitor of TNF-α production from RBL-2H3 cells. Biol. Pharm. Bull. 31(5), 831-833 (2008).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website

Copyright Cayman Chemical Company, 02/25/2022

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM