



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 


PRODUCT INFORMATION



L-threo-Sphingosine (d18:1)

Item No. 10010541

CAS Registry No.: 25695-95-8
Formal Name: 2S-amino-4E-octadecene-1,3S-diol
Synonym: L-threo-Sphingosine C18
MF: C₁₈H₃₇NO₂
FW: 299.5
Purity: ≥98%
Supplied as: A crystalline solid
Storage: -20°C
Stability: ≥2 years



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

L-threo-Sphingosine (d18:1) is supplied as a crystalline solid. A stock solution may be made by dissolving the L-threo-sphingosine (d18:1) in the solvent of choice. L-threo-Sphingosine (d18:1) is soluble in the organic solvent ethanol, which should be purged with an inert gas, at a concentration of approximately 0.25 mg/ml.

Description

Sphingosines are long-chain base precursors of cellular sphingolipids and are used directly in the synthesis of ceramide. Sphingosine can exist in four stereoisomers, however only D-erythro-sphingosine occurs naturally. L-threo-Sphingosine C-18, an analog of D-erythro-sphingosine, inhibits protein kinase C in mixed micelle assays with 50% inhibition at 2.2 mol % making it a slightly more potent inhibitor compared to D-erythro-sphingosine (50% inhibition at 2.8 mol %) or other analogs of shorter alkyl chain length.¹ Addition of L-threo-Sphingosine C-18 to primary cultured cerebellar cells does not decrease serine palmitoyltransferase (SPT) activity, the rate-limiting enzyme in ceramide biosynthesis, whereas under similar conditions D-erythro-sphingosine inhibits SPT activity.²

References

1. Merrill, A.H., Jr., Nimkar, S., Menaldino, D., *et al.* Structural requirements for long-chain (shingoid) base inhibition of protein kinase C *in vitro* and for the cellular effects of these compounds. *Biochemistry* **28**(8), 3138-3145 (1989).
2. Mandon, E.C., van Echten, G., Birk, R., *et al.* Shingolipid biosynthesis in cultured neurons: Down-regulation of serine palmitoyltransferase by sphingoid bases. *Eur. J. Biochem.* **198**(3), 667-674 (1991).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the [complete](#) Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 02/06/2018

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897
[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM