

Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com

PRODUCT INFORMATION



Digoxin-d₃ Item No. 10010657

CAS Registry No.:		
Formal Name:	(3β,5β,12β)-3-[(O-2,6-dideoxy-β-D-ribo-	0
	hexopyranosyl-(1→4)-O-2,6-dideoxy-β-D-	A
	ribo-hexopyranosyl- $(1\rightarrow 4)$ -2,6-dideoxy- β -D-	D-{\ Ĭ
	ribo-hexopyranosyl)oxy]-12,14-dihydroxy-	он С
	card-20(22)-enolide-21,21,22-d ₃	
MF:	C ₄₁ H ₆₁ D ₃ O ₁₄	
FW:	784.0	
Chemical Purity:	≥95% (Digoxin)	
Deuterium		
Incorporation:	≥99% deuterated forms (d₁-d₃); ≤1% d₀	
Supplied as:	A solid	1
Storage:	-20°C	
Stability:	≥2 years	
Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.		

Laboratory Procedures

Digoxin-d₂ is intended for use as an internal standard for the quantification of digoxin (Item No. 22266) by GC- or LC-MS. The accuracy of the sample weight in this vial is between 5% over and 2% under the amount shown on the vial. If better precision is required, the deuterated standard should be quantitated against a more precisely weighed unlabeled standard by constructing a standard curve of peak intensity ratios (deuterated versus unlabeled).

Digoxin-d₃ is supplied as a solid. A stock solution may be made by dissolving the digoxin-d₃ in the solvent of choice, which should be purged with an inert gas. Digoxin- d_3 is soluble in organic solvents such as methanol (warmed) and DMSO.

Description

Digoxin is a cardiac glycoside and metabolite of digitoxin (Item No. 27825) that binds to and inhibits the Na⁺/K⁺-ATPase in cardiac tissues in an ATP- and Mg^{2+} -dependent manner.¹ This inhibition results in loss of the transmembrane Na⁺ gradient, which decreases activity of the Na⁺/Ca²⁺ exchanger, increasing intracellular Ca²⁺ levels, inotropy, and cardiac force.² It increases activity of mitochondrial ATPase and actomyosin ATPase in rat hearts, which is directly correlated with increased myofibrillar contractile strength.³ In vivo, digoxin also decreases right atrial pressure and increases cardiac output in a canine model of congestive heart failure produced by pulmonary artery constriction.⁴ Formulations containing digoxin have been used to treat atrial fibrillation.⁵

References

- 1. Matsui, H. and Schwartz, A. Biochim. Biophys. Acta. 151(3), 655-663 (1968).
- 2. Neves, C.H., Tibana, R.A., Prestes, J., et al. Int. J. Sports Med. 38(4), 263-269 (2017).
- 3. Hamrick, M.E. and Fritz, P.J. Biochem. Biophys. Res. Commun. 22(5), 540-546 (1966).
- 4. Davis, J.O., Howell, D.S., and Hyatt, R.E. Circ. Res. 3(3), 259-263 (1955).
- 5. Kotecha, D., Calvert, M., Deeks, J.J., et al. BMJ Open 7(7), e015099 (2017).

WARNING THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFFTY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

uyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 06/26/2020

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897 [734] 971-3335 FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.CAYMANCHEM.COM