

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



Chenodeoxycholic Acid

Item No. 10011286

CAS Registry No.: 474-25-9

Formal Name: $(3\alpha,5\beta,7\alpha)$ -3,7-dihydroxy-cholan-24-oic acid Synonyms: Anthropodeoxycholic Acid, CDCA, Fluibil,

Hekbilin, Kebilis, Ulmenide

MF: $C_{24}H_{40}O_4$ FW: 392.6 **Purity:** ≥95%

Supplied as: A crystalline solid

Storage: -20°C Stability: ≥2 vears

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

Chenodeoxycholic acid (CDCA) is supplied as a crystalline solid. A stock solution may be made by dissolving the CDCA in an organic solvent purged with an inert gas. CDCA is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide (DMF). The solubility of CDCA in ethanol and DMSO is approximately 20 mg/ml and approximately 30 mg/ml in DMF.

CDCA is sparingly soluble in aqueous buffers. For maximum solubility in aqueous buffers, CDCA should first be dissolved in DMF and then diluted with the aqueous buffer of choice. CDCA has a solubility of approximately 0.5 mg/ml in a 1:1 solution of DMF:PBS (pH 7.2) using this method. We do not recommend storing the aqueous solution for more than one day.

Description

CDCA is a hydrophobic primary bile acid. It is formed from cholesterol in the liver via a multistep process catalyzed by the cytochrome P450 (CYP) isoforms CYP7A1, CYP8B1, and CYP27A1. CDCA is a farnesoid X receptor (FXR) agonist that binds to FXRs in a TR-FRET assay (EC $_{50}$ = 13 μ M) and induces FXR transactivation in a reporter assay. $^{2.3}$ It induces transcription of the gene encoding the Nrf2 target glutamate cysteine ligase (GCL) in primary hepatocytes and HepG2 cells when used at concentrations ranging from 25 to 100 μΜ.⁴

References

- 1. Fiorucci, S. and Distrutti, E. Chenodeoxycholic acid: An update on its therapeutic applications. Bile acids and their receptors. Handbook of experimental pharmacology. Fiorucci, S. and Distrutti, E., editors, 1st edition, Springer (2019).
- 2. Ohinata, Y., Payer, B., O'Carroll, D., et al. Blimp1 is a critical determinant of the germ cell lineage in mice. Nature 436, 207-213 (2005).
- Urizar, N.L., Liverman, A.B., Dodds, D.T., et al. A natural product that lowers cholesterol as an anatagonist ligand for FXR. Science 296(5573), 1703-1706 (2002).
- Tan, K.P., Yang, M., and Ito, S. Activation of nuclear factor (erythroid-2 like) factor 2 by toxic bile acids provokes adaptive defense responses to enhance cell survival at the emergence of oxidative stress. Mol. Pharmacol. 72(5), 1380-1390 (2007).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website

Copyright Cayman Chemical Company, 09/28/2020

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM