

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



N-Oleoyl Dopamine

Item No. 10115

CAS Registry No.: 105955-11-1

N-[2-(3,4-dihydroxyphenyl)ethyl]-Formal Name:

9Z-octadecenamide

Synonym: MF: $C_{26}H_{43}NO_3$ FW: 417.6 ≥98% **Purity:**

≥1 year at -20°C Stability: Supplied as: A crystalline solid λ_{max} : 283 nm UV/Vis.:

Laboratory Procedures

For long term storage, we suggest that N-oleoyl dopamine (ODA) be stored as supplied at -20°C. It should be stable for at least one year.

ODA is supplied as a crystalline solid. A stock solution may be made by dissolving the ODA in the solvent of choice. ODA is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide, which should be purged with an inert gas. The solubility of ODA in these solvents is approximately 50, 20, and 30 mg/ml, respectively.

ODA is sparingly soluble in aqueous buffers. For maximum solubility in aqueous buffers, ODA should first be dissolved in ethanol and then diluted with the aqueous buffer of choice. ODA has a solubility of approximately 500 μg/ml in a 1:1 solution of ethanol:PBS (pH 7.2) using this method. We do not recommend storing the aqueous solution for more than one day.

Description

ODA is a selective, endogenous vanilloid receptor 1 (VR₁) agonist isolated from bovine brain. ¹ Structurally, it is the amide of oleic acid and dopamine and is therefore a "hybrid" analog which incorporates components of both the anadamide-like and dopamine neurotransmitter pathways. ODA binds to the human recombinant VR₁ with a K_i of 36 nM making it equipotent to capsaicin and slightly more potent than N-arachidonoyl dopamine in this assay. It causes hyperalgesia and nocifensive behavior that is blocked by the VR₁ antagonist iodo-resiniferatoxin. ODA is selective for VR₁ based on observations that it has weak affinity for the rat CB₁ receptor (K, of 1.6 μM) and is a very weak inhibitor of FAAH. ODA is also a potent inhibitor of 5-lipoxygenase from rat basophilic leukemia-1 (RBL-1) cells, with a IC₅₀ of 7.5 nM.^{2,3}

References

- 1. Chu, C.J., Huang, S.M., De Petrocellis, L., et al. N-oleoyldopamine, a novel endogenous capsaicin-like lipid that produces hyperalgesia. J. Biol. Chem. 278(16), 13633-13639 (2003).
- Tseng, C.-F., Iwakami, S., Mikajiri, A., et al. Inhibition of in vitro prostaglandin and leukotriene biosyntheses by cinnamoyl-β-phenethylamine and N-acyldopamine derivatives. Chem. Pharm. Bull. 40(2), 396-400
- 3. Iwakami, S., Shibuya, M., Tseng, C.-F., et al. Inhibition of arachidonate 5-lipoxygenase by phenolic compounds. Chem. Pharm. Bull. 34, 3960-3963 (1986).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

uyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website

Copyright Cayman Chemical Company, 08/10/2015

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM