

Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com

PRODUCT INFORMATION



COOH

13-epi-12-oxo Phytodienoic Acid

Item No. 10195

CAS Registry No.:	71606-07-0	
Formal Name:	4-oxo-5R-(2Z)-2-pentenyl-2-	
	cylcopentene-1S-octanoic acid	0
Synonym:	13-epi-12-oxo PDA	
MF:	C ₁₈ H ₂₈ O ₃	
FW:	292.4	
Purity:	≥95%	
Stability:	≥1 year at -80°C	• ~
Supplied as:	A solution in ethanol	
UV/Vis.:	λ _{max} : 220 nm	

Laboratory Procedures

For long term storage, we suggest that 13-epi-12-oxo phytodeinoic acid (13-epi-12-oxo PDA) be stored as supplied at -80°C. It should be stable for at least one year.

13-epi-12-oxo PDA is supplied as a solution in ethanol. To change the solvent, simply evaporate the ethanol under a gentle stream of nitrogen and immediately add the solvent of choice. Solvents such as DMSO and dimethyl formamide purged with an inert gas can be used. The solubility of 13-epi-12-oxo PDA in these solvents is approximately 10 and 20 mg/ml, respectively.

Further dilutions of the stock solution into aqueous buffers or isotonic saline should be made prior to performing biological experiments. Ensure that the residual amount of organic solvent is insignificant, since organic solvents may have physiological effects at low concentrations. If an organic solvent-free solution of 13-epi-12-oxo PDA is needed, it can be prepared by evaporating the ethanol and directly dissolving the neat oil in aqueous buffers. The solubility of 13-epi-12-oxo PDA in PBS (pH 7.2) is approximately 1 mg/ml. We do not recommend storing the aqueous solution for more than one day.

Description

13-epi-12-oxo PDA is a lipoxygenase metabolite of α -linolenic acid in the leaves of green plants such as corn. ω -3 and ω -6 polyunsaturated fatty acids in plants are substrates for plant lipoxygenases.¹ 12-oxo PDA is one of the best studied end metabolites of this enzymatic pathway.² While the initial enzymatic product and major isomer of 12-oxo PDA contains side chains in the cis position, both being β to the ring, the upper side chain attached at C-13 can and frequently does isomerize when 12-oxo PDA is extracted, isolated or stored. 13-epi-12-oxo PDA is the product of this isomerization.

References

- 1. Crombie, L. and Mistry, K.M. Synthesis of 12-oxophytodienoic acid (12-oxo PDA) and the compounds of its enzymic degradation cascade in plants, OPC-8:0, -6:0, -4:0 and -2:0 (epi-jasmonic acid), as their methyl esters. J. Chem. Soc. 1, 1981-1991 (1991).
- 2. Hamberg, M. and Gardner, H.W. Oxylipin pathway to jasmonates: Biochemistry and biological significance. Biochim. Biophys. Acta 1165, 1-18 (1992).

WARNING THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFFTY DATA

al should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution

WARRANTY AND LIMITATION OF REMEDY

uyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 08/12/2015

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA **PHONE:** [800] 364-9897 [734] 971-3335 FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.CAYMANCHEM.COM