



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



PRODUCT INFORMATION




CD3 Monoclonal FITC Antibody (Clone RIV9)

Item No. 10231

Overview and Properties

Contents:	This vial contains 100 µg of fluorescein-labeled affinity-purified monoclonal antibody.
Cross Reactivity:	(+) Human
Form:	Liquid
Storage:	-20°C (as supplied)
Stability:	≥1 year
Storage Buffer:	500 µl PBS, pH 7.2, with 50% glycerol, 0.1% BSA, and 0.02% sodium azide
Clone:	RIV9
Host:	Mouse
Isotype:	IgG3
Applications:	Flow cytometry (FC), immunofluorescence (IF), and immunohistochemistry (IHC) (frozen sections); the recommended starting dilution for FC is 5 µg/ml (25 µl/test) and 10-20 µg/ml for IF and IHC (frozen sections). Not recommended for IHC (paraffin-embedded tissue) or Western blot. Other applications were not tested, therefore optimal working concentration/dilution should be determined empirically.

Image



WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 07/10/2023

CAYMAN CHEMICAL
1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 - USA
PHONE: [800] 364-9897
[734] 971-3335
FAX: [734] 971-3640
CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM

PRODUCT INFORMATION

Description

The CD3 antigen is a specific T-cell marker expressed on thymocytes and mature T-lymphocytes. The CD3 complex contains CD3g ($M_r = 26$ kDa), CD3d ($M_r = 21$ kDa), and CD3e ($M_r = 20$ kDa).¹ CD3 is detected on the surface membrane or in the cytoplasm of T-cells from peripheral blood, depending on the state of cell differentiation. In addition to detection of T-cells from peripheral blood, the CD3 antibody can be used for detection of reactive T-cells in lymphoid infiltrates in tissues and neoplastic T-cells in non-Hodgkin's lymphomas of T-cell lineage.^{2,3}

References

1. Berkhouit, B., Alarcon, B., and Terhorst, C. Transfection of genes encoding the T-cell receptor-associated CD3 complex into COS cells results in assembly of the macromolecular structure. *J. Biol. Chem.* **263**(17), 8528-8536 (1988).
2. Ceuppens, J.L. and Van Vaeck, F. Human T cell activation induced by a monoclonal mouse IgG₃ anti-CD3 antibody (RIV9) requires binding of the Fc part of the antibody to the monocytic 72-kDa high-affinity Fc receptor (FcRI)1. *Cell. Immunol.* **118**, 136-146 (1989).
3. Jiskoot, W., Hoven, A.-M.V., De Koning, A.A.M., et al. Purification and stabilisation of a poorly soluble mouse IgG₃ monoclonal antibody. *J. Immunol. Methods* **138**, 181-189 (1991).