



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



PRODUCT INFORMATION




PCSK9 (human) Polyclonal Antibody

Item No. 10240

Overview and Properties

Contents:	This vial contains 500 µg of protein A-purified polyclonal antibody.
Synonyms:	NARC-1, Proprotein Convertase Subtilisin Kexin 9
Immunogen:	Purified recombinant human PCSK9
Species Reactivity:	(+) Human, mouse, and rat
Uniprot No.:	Q8NBP7
Form:	Liquid
Storage:	-20°C (as supplied)
Stability:	≥3 years
Storage Buffer:	PBS, pH 7.2, with 50% glycerol and 0.02% sodium azide
Host:	Rabbit
Application:	Western blot (WB); the recommended starting dilution is 1:200. Other applications were not tested, therefore optimal working concentration/dilution should be determined empirically.

Image



Lane 1: Mouse heart (10,000 x g supernatant) (40 µg)

Lane 2: Mouse liver (10,000 x g supernatant) (50 µg)

Lane 3: PCSK9 Western Ready Control (2 µl)

Lane 4: PCSK9 Western Ready Control (5 µl)

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 11/03/2023

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 - USA

PHONE: [800] 364-9897
[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM

PRODUCT INFORMATION

Description

PCSK9 is a member of the subtilisin serine protease family with an important role in low-density lipoprotein (LDL) metabolism.¹ Mutation in the PCSK9 gene is associated with autosomal dominant hypercholesterolemia which is characterized by an increase in LDL cholesterol levels.² PCSK9 overexpression in wild-type mice doubles the plasma total cholesterol, possibly through acceleration of the degradation of the LDL receptor.^{1,3} PCSK9 mRNA is detected in various tissues such as liver, kidney, lung, spleen, jejunum, ileum, colon, and muscles with the highest expression in the liver.⁴ Human PCSK9 precursor is 692 amino acids in length with an estimated molecular weight of 74 kDa. This proprotein is self-cleaved to form a mature protein of 63 kDa in the Golgi.⁵

References

1. Maxwell, K.N., Fisher, E.A., and Breslow, J.L. Overexpression of PCSK9 accelerates the degradation of the LDLR in a post-endoplasmic reticulum compartment. *Proc. Natl. Acad. Sci. USA* **102**(6), 2069-2074 (2005).
2. Abifadel, M., Varret, M., Rabès, J.-P., et al. Mutations in PCSK9 cause autosomal dominant hypercholesterolemia. *Nature Genet.* **34**(2), 154-156 (2003).
3. Maxwell, K.N. and Breslow, J.L. Adenoviral-mediated expression of PCSK9 in mice results in a low-density lipoprotein receptor knockout phenotype. *Proc. Natl. Acad. Sci. USA* **101**(18), 7100-7105 (2004).
4. Seidah, N.G., Benjannet, S., Wickham, L., et al. The secretory proprotein convertase neural apoptosis-regulated convertase 1 (NARC-1): Liver regeneration and neuronal differentiation. *Proc. Natl. Acad. Sci. USA* **100**(3), 928-933 (2003).
5. Maxwell, K.N. and Breslow, J.L. Proprotein convertase subtilisin kexin 9: The third locus implicated in autosomal dominant hypercholesterolemia. *Curr. Opin. Lipidol.* **16**, 167-172 (2005).