

# Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



## Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

## Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

## SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



# PRODUCT INFORMATION



## 16-phenoxy tetranor Prostaglandin A<sub>2</sub>

Item No. 10285

CAS Registry No: 51639-10-2

Formal Name: 9-oxo-15R-hydroxy-16-phenoxy-

17,18,19,20-tetranor-prosta-

5Z,10,13E-trien-1-oic acid


Synonym: 16-phenoxy tetranor PGA<sub>2</sub>

MF:  $C_{22}H_{26}O_5$ FW: 370.4 **Purity:** ≥98%

Stability: ≥1 year at -20°C

Supplied as: A solution in methyl acetate

UV/Vis:  $\lambda_{max}$ : 219 nm



### **Laboratory Procedures**

For long term storage, we suggest that 16-phenoxy tetranor prostaglandin  $A_2$  (16-phenoxy tetranor PGA<sub>2</sub>) be stored as supplied at -20°C. It should be stable for at least one year.

16-phenoxy tetranor PGA2 is supplied as a solution in methyl acetate. To change the solvent, simply evaporate the methyl acetate under a gentle stream of nitrogen and immediately add the solvent of choice. Solvents such as DMSO, ethanol, and dimethyl formamide purged with an inert gas can be used. The solubility of 16-phenoxy tetranor  $PGA_2$  in these solvents is at least 100 mg/ml. 16-phenoxy tetranor  $PGA_2$ is stable for at least six months in these solvents if stored at -20°C.

Further dilutions of the stock solution into aqueous buffers or isotonic saline should be made prior to performing biological experiments. Ensure that the residual amount of organic solvent is insignificant since organic solvents may have physiological effects at low concentrations. If an organic solvent-free solution of 16-phenoxy tetranor PGA2 is needed, it can be prepared by evaporating the methyl acetate and directly dissolving the neat oil in aqueous buffers. The solubility of 16-phenoxy tetranor PGA<sub>2</sub> in PBS (pH 7.2) is at least 0.8 mg/ml. Store aqueous solutions of 16-phenoxy tetranor PGA2 on ice and use within 12 hours of preparation.

#### Description

16-phenoxy tetranor PGA<sub>2</sub> is a minor metabolite found in human plasma after intravenous administration of sulprostone. Its biological activity has not been studied or reported in the literature.

#### Reference

1. Kuhnz, W., Hoyer, G., Backhus, S., et al. Identification of the major metabolites of the prostaglandin E<sub>2</sub>-analogue sulprostone in human plasma, and isolation from urine (in vivo) and liver perfusate (in vitro) of female guinea-pigs. Drug Metab. Dispos. 19, 920-925 (1991).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

## WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 08/13/2015

### **CAYMAN CHEMICAL**

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

**FAX:** [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM