

Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com

PRODUCT INFORMATION



8,12-*iso*-iPF_{2 α}-VI-1,5-lactone Item No. 10312

Formal Name:	6-((E)-2-((1R,2S,3R,5S)-3,5- dihydroxy-2-((Z)-oct-2-enyl) cyclopentyl)vinyl)tetrahydro-2H- pyran-2-one	но
MF:	C ₂₀ H ₃₂ O ₄	
FW:	336.5	
Purity:	≥98%	
Stability:	≥1 year at -80°C	
Supplied as:	A solution in methyl acetate	нÓ

Laboratory Procedures

For long term storage, we suggest that 8,12-iso-iPF $_{2n}$ -VI-1,5- lactone be stored as supplied at -80°C. It should be stable for at least one year.

8,12-iso-iPF_{2a}-VI-1,5- lactone is supplied as a solution in methyl acetate. To change the solvent, simply evaporate the methyl acetate under a gentle stream of nitrogen and immediately add the solvent of choice. Solvents such as ethanol, DMSO, and dimethyl formamide purged with an inert gas can be used. The solubility of 8,12-iso-iPF_{2a}-VI-1,5- lactone in these solvents is approximately 100 mg/ml.

Further dilutions of the stock solution into aqueous buffers or isotonic saline should be made prior to performing biological experiments. Ensure that the residual amount of organic solvent is insignificant, since organic solvents may have physiological effects at low concentrations. If an organic solvent-free solution of 8,12-*iso*-iPF_{2α}-VI-1,5- lactone is needed, it can be prepared by evaporating the methyl acetate and directly dissolving the neat oil in aqueous buffers. The solubility of 8,12-*iso*-iPF_{2α}-VI-1,5- lactone in PBS, pH 7.2, is approximately 1 mg/ml. We do not recommend storing the aqueous solution for more than one day.

Description

F₂ isoprostanes (F₂-iPs) are thought to arise from the free radical-mediated peroxidation of phospholipidbound arachidonic acid.¹ They are cleaved, presumably by an unidentified phospholipase A_2 , and are found in the circulation and the urine. 8,12-iso-iPF_{2a}-VI-1,5-lactone is a racemic mixture of the lactone form of the free acid, 8,12-iso- iPF_{2a}-VI. Previously called IPF_{2a}-I, the free acid form, iPF_{2a}-VI, is the most abundant F_2 -iP regioisomer measured in the urine of rats treated with CCl₄ to induce lipid peroxidation.^{2,3} iPF_{2a}-VI is the only regioisomer that undergoes lactonization, and this occurs slowly in vivo or can be driven chemically.⁴ The less polar lactone is readily separated from the free acid forms of iPF_{2a} .⁴ While the level of iPF_{2a} -VI in plasma, urine, and organs is used as a biomarker for oxidative stress, some F_2 -iPs also evoke significant biological effects.⁵⁻⁷ It is not known if 8,12-*iso*-iPF_{2a}-VI-1,5-lactone has important physiological effects.

References

- 1. Lawson, J.A., Li, H., Rokach, J., et al. J. Biol. Chem. 273, 29295-29301 (1998).
- 2. Morrow, J.D., Hill, K.E., Burk, R.F., et al. Proc. Natl. Acad. Sci. USA 87, 9383-9387 (1990).
- 3. Waugh, R.J., Morrow, J.D., Roberts, L.J., II, et al. Free Radic. Biol. Med. 23, 943-954 (1997).
- 4. Adiyaman, M., Lawson, J.A., Khanapure, S.P., et al. Anal. Biochem. 262, 45-56 (1998).
- 5. Practico, D., Barry, O.P., Lawson, J.A., et al. Proc. Natl. Acad. Sci. USA 95, 3449-3454 (1998).
- 6. Van Eck, M., Hoekstra, M., Hildebrand, R.B., et al. Arterioscler. Thromb. Vasc. Biol. 27, 2413-2419 (2007).
- 7. Doe, C., Bentley, R., Behm, D.J., et al. J. Pharmacol. Exp. Ther. 320(1), 89-98 (2007).

WARNING THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution

WARRANTY AND LIMITATION OF REMEDY

uver agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website

Copyright Cayman Chemical Company, 08/19/2015

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897 [734] 971-3335 FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.CAYMANCHEM.COM