



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# PRODUCT INFORMATION




## TIM-4 Monoclonal Antibody

Item No. 10322

### Overview and Properties

<b>Contents:</b>	This vial contains 400 µg of protein A-purified monoclonal antibody.
<b>Synonyms:</b>	T cell Immunoglobulin and Mucin Domain-containing 4, T Cell Immunoglobulin Mucin Receptor 4, T cell Membrane Protein 4, TIMD-4
<b>Immunogen:</b>	Mouse peritoneal cell TIM-4
<b>Species Reactivity:</b>	(+) Human; other species not tested
<b>Cross Reactivity:</b>	(+) TIM-4
<b>Uniprot No.:</b>	Q96H15
<b>Form:</b>	Liquid
<b>Storage:</b>	-20°C (as supplied)
<b>Stability:</b>	≥3 years
<b>Storage Buffer:</b>	PBS, pH 7.2, with 50% glycerol and 0.02% sodium azide
<b>Host:</b>	Armenian hamster
<b>Applications:</b>	Flow cytometry (FC) and immunocytochemistry (ICC); The optimal working concentration/dilution should be determined empirically.

### Image



Black: Blank  
Red: Normal Rabbit IgG-FITC (0.01 µg/ml)  
Blue: TIM-4 Monoclonal Antibody (1 µg/ml)  
Yellow: TIM-4 Monoclonal Antibody (5 µg/ml)  
Green: TIM-4 Monoclonal Antibody (10 µg/ml)

Jurkat cells were fixed with cytospin, blocked with 5% normal goat serum, and washed between steps.  
2°Ab used = Goat anti-hamster-FITC: 1:750.

WARNING  
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA  
This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY  
Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 11/03/2023

CAYMAN CHEMICAL  
1180 EAST ELLSWORTH RD  
ANN ARBOR, MI 48108 · USA  
PHONE: [800] 364-9897  
[734] 971-3335  
FAX: [734] 971-3640  
CUSTSERV@CAYMANCHEM.COM  
WWW.CAYMANCHEM.COM

# PRODUCT INFORMATION



## Description

---

TIM-4 is a type I transmembrane protein and member of the T cell immunoglobulin and mucin domain-containing (TIM) family of immunoregulatory proteins.<sup>1</sup> It is composed of an N-terminal immunoglobulin variable (IgV) domain that binds to phosphatidylserine, as well as a mucin stalk, which contains O- and N-linked glycosylation sites, a transmembrane domain, and a C-terminal cytoplasmic tail, which lacks a tyrosine phosphorylation site, unlike TIM-1 and TIM-3.<sup>1,2</sup> TIM-4 is expressed in antigen-presenting cells (APCs) such as dendritic cells and macrophages and binds to TIM-1 on activated T cells.<sup>3</sup> It is mainly involved in phagocytosis of apoptotic cells *via* recognition of phosphatidylserine, but also plays a role in T cell proliferation and survival, viral entry, and antitumor immunity.<sup>2,4,5,6</sup> TIM-4 targeting antibodies prevent HIV-1 entry into host cells *in vitro* and enhance the efficacy of anticancer vaccines in a murine melanoma model.<sup>5,6</sup> SNPs in *TIMD4* are associated with increased susceptibility of asthma in children.<sup>7</sup> Cayman's TIM-4 Monoclonal Antibody can be used for flow cytometry (FC) and immunocytochemistry (ICC) applications.

## References

---

1. Freeman, G.J., Casasnovas, J.M., Umetsu, D.T., *et al.* TIM genes: A family of cell surface phosphatidylserine receptors that regulate innate and adaptive immunity. *Immunol. Rev.* **235(1)**, 172-189 (2010).
2. Miyanishi, M., Tada, K., Koike, M., *et al.* Identification of Tim4 as a phosphatidylserine receptor. *Nature* **450**, 435-439 (2007).
3. Meyers, J.H., Chakravarti, S., Schlesinger, D., *et al.* TIM-4 is the ligand for TIM-1, and the TIM-1-TIM-4 interaction regulates T cell proliferation. *Nat. Immunol.* **6**, 455-464 (2005).
4. Rodriguez-Manzanet, R., Meyers, J.H., Balasubramanian, S., *et al.* TIM-4 expressed on APCs induces T cell expansion and survival. *J. Immunol.* **180(7)**, 4706-4713 (2008).
5. Sims, B., Farrow, A.L., Williams, S.D., *et al.* Role of TIM-4 in exosome-dependent entry of HIV-1 into human immune cells. *Int. J. Nanomedicine* **12**, 4823-4833 (2017).
6. Baghdadi, M., Nagao, H., Yoshiyama, H., *et al.* Combined blockade of TIM-3 and TIM-4 augments cancer vaccine efficacy against established melanomas. *Cancer Immunol. Immunother.* **62(4)**, 629-637 (2013).
7. Cai, P.C., Hu, L.H., Cui, T.P., *et al.* Association of TIM4 promoter polymorphism -1419G>A with childhood asthma in a Chinese Han population. *Tissue Antigens* **74**, 11-16 (2009).

CAYMAN CHEMICAL  
1180 EAST ELLSWORTH RD  
ANN ARBOR, MI 48108 · USA  
PHONE: [800] 364-9897  
[734] 971-3335  
FAX: [734] 971-3640  
CUSTSERV@CAYMANCHEM.COM  
WWW.CAYMANCHEM.COM