

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



Cytokeratin Monoclonal PE Antibody (Clone C-11)

Item No. 10478

Overview and Properties

This vial contains 100 µg of protein A-purified, PE-labeled monoclonal antibody. Contents:

Synonyms: CKs, pan-Cytokeratin, Keratin

Cross Reactivity: (+) Cytokeratins 4, 5, 6, 8, 10, 13, and 18

Species Reactivity: (+) Human Form: Lyophilized -20°C (as supplied) Storage:

Stability: ≥1 year

Storage Buffer: 100 mM sodium phosphate, pH 7.4, with 50 mM sucrose, 150 mM sodium chloride,

0.1% BSA, and 2 mM sodium azide when reconstituted in 1 ml double distilled water

Clone: Mouse Host: Isotype: lgG1


Flow cytometry (FC) and Immunofluorescence (IF). The recommended starting dilution **Applications:**

for FC and IF is 5-10 µg/ml. Other applications were not tested, therefore optimal

working concentration/dilution should be determined empirically.

Miscellaneous: Avoid exposure to heat and light

Image



HeLa cells stained with Cytokeratin Monoclonal PE Antibody (Clone C-11), 10 μg/ml.

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website

Copyright Cayman Chemical Company, 10/23/2023

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.CAYMANCHEM.COM

PRODUCT INFORMATION



Description

Cytokeratins (CKs) are intermediate filaments proteins and members of the cytokeratin protein family. $^{1-3}$ The cytokeratin family is composed of at least 20 unique gene products, which fall into two categories: type I acidic cytokeratins, CK9-CK20, and type II neutral-basic cytokeratins, CK1-CK8. Each cytokeratin is composed of a central rod containing four α -helical domains, which are important for self-assembly, and non-helical head and tail domains at the N- and C-termini, respectively. Cytokeratins are major structural proteins expressed in epithelial cells and localized to the cytoplasm where they form heterozygous type I-type II pairs. They primarily function to protect epithelial cells from mechanical and non-mechanical stresses that induce cell rupture and death. Cytokeratins have additional isoform- and tissue-specific functions, including the modulation of protein synthesis and cell size during epithelial growth and roles in skin pigmentation. Cayman's Cytokeratin Monoclonal PE Antibody (Clone C-11) can be used for flow cytometry (FC) and immunofluorescence (IF) applications. This antibody recognizes cytokeratins 4, 5, 6, 8, 10, 13, and 18 from human samples.

References

- Weng, Y.-R., Cui, Y., and Fang, J.-Y. Biological functions of cytokeratin 18 in cancer. Mol. Cancer Res. 10(4), 485-493 (2012).
- 2. 15723Gu, L.-H. and Coulombe, P.A. Keratin function in skin epithelia: A broadening palette with surprising shades. *Curr. Opin. Cell Biol.* 19(1), 13-23 (2007).
- 3. Eldirany, S.A., Lomakin, I.B., Ho, M., et al. Recent insight into intermediate filament structure. Curr. Opin. Cell Biol. 68, 132-143 (2021).
- 4. Ditzel, H.J., Strik, M.C.M., Larsen, M.K., *et al.* Cancer-associated cleavage of cytokeratin 8/18 heterotypic complexes exposes a neoepitope in human adenocarcinomas. *J. Biol. Chem.* **277(24)**, 21712-21722 (2002).

ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897