



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com


[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

PRODUCT INFORMATION



5(Z),8(Z),11(Z)-Eicosatrienoic Acid-d₆ Item No. 10742

Formal Name: 5Z,8Z,11Z-eicosatrienoic-5,6,8,9,11,12-d₆ acid
Synonym: Mead Acid-d₆
MF: C₂₀H₂₈D₆O₂
FW: 312.5
Chemical Purity: ≥98% (5(Z),8(Z),11(Z)-Eicosatrienoic Acid)
Deuterium Incorporation: ≥99% deuterated forms (d₁-d₆); ≤1% d₀
Supplied as: A solution in methyl acetate
Storage: -20°C
Stability: ≥1 year



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

5(Z),8(Z),11(Z)-Eicosatrienoic acid-d₆ (mead acid-d₆) is intended for use as an internal standard for the quantification of mead acid (Item No. 90190) by GC- or LC-MS. The accuracy of the sample weight in this vial is between 5% over and 2% under the amount shown on the vial. If better precision is required, the deuterated standard should be quantitated against a more precisely weighed unlabeled standard by constructing a standard curve of peak intensity ratios (deuterated versus unlabeled).

Mead acid-d₆ is supplied as a solution in methyl acetate. To change the solvent, simply evaporate the methyl acetate under a gentle stream of nitrogen and immediately add the solvent of choice. Solvents such as DMSO and dimethyl formamide purged with an inert gas can be used. The solubility of mead acid-d₆ in these solvents is approximately 100 mg/ml.

Description

Mead acid accumulates in the tissues of animals fed diets deficient in both ω-3 and ω-6 fatty acids. It can be converted to 3-series cysteinyl-leukotrienes but cannot serve as a COX substrate.¹ The kidneys from essential fatty acid deficient rats are less immunogenic when transplanted, and the metabolites of eicosatrienoic acid are believed to play a role in altering the immune status of these organs.²

References

1. Lefkowitz, J.B. Essential fatty acid deficiency: Probing the role of arachidonate in biology. *Adv. Prostaglandin Thromboxane Leukotriene Res.* **20**, 224-231 (1990).
2. Schreiner, G.F., Flye, W., Brunt, E., *et al.* Essential fatty acid depletion of renal allografts and prevention of rejection. *Science* **240**, 1032-1033 (1988).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 07/06/2018

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897
[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM