

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



Everolimus

Item No. 11597

CAS Registry No.: 159351-69-6


42-O-(2-hydroxyethyl)-rapamycin Formal Name:

RAD001 Synonym: MF: C₅₃H₈₃NO₁₄ FW: 958.2 **Purity:**

 λ_{max} : 268, 277, 289 nm A crystalline solid UV/Vis.: Supplied as:

-20°C Storage: Stability: ≥4 years

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.



Laboratory Procedures

Everolimus is supplied as a crystalline solid. A stock solution may be made by dissolving the everolimus in the solvent of choice, which should be purged with an inert gas. Everolimus is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide (DMF). The solubility of everolimus in ethanol and DMSO is approximately 10 mg/ml and approximately 20 mg/ml in DMF.

Everolimus is sparingly soluble in aqueous buffers. For maximum solubility in aqueous buffers, everolimus should first be dissolved in DMF and then diluted with the aqueous buffer of choice. Everolimus has a solubility of approximately 0.1 mg/ml in a 1:4 solution of DMF:PBS (pH 7.2) using this method. We do not recommend storing the aqueous solution for more than one day.

Description

The mammalian target of rapamycin (mTOR) is a serine/threonine kinase that, as part of two distinct complexes (mTORC1 and mTORC2), plays pivotal roles in intracellular signaling. 1-3 Everolimus is a hydroxyethyl ether rapamycin (Item No. 13346) derivative that inhibits mTOR signaling through both mTORC1 and mTORC2 when added to cells at 20 nM.^{4,5} It is orally available and shows improved pharmacokinetics and pharmacodynamics over rapamycin.⁵ Through its inhibition of mTOR, everolimus inhibits cell proliferation, metabolism, and angiogenesis in certain types of cancer.^{5,6} It also acts as an immunosuppressive agent in the context of organ transplantation.^{5,7}

References

- 1. Petroulakis, E., Mamane, Y., Le Bacquer, O., et al. Br. J. Cancer 94(2), 195-199 (2006).
- 2. Dann, S.G., Selvaraj, A., and Thomas, G. Trends Mol. Med. 13(6), 252-259 (2007).
- 3. Tao, Z., Barker, J., Shi, S.D.H., et al. Biochem. 49(39), 8488-8498 (2010).
- 4. Zeng, Z., Sarbassov, D.D., Samudio, I.J., et al. Blood 109(8), 3509-3512 (2007).
- 5. Lebwohl, D., Anak, Ö., Sahmoud, T., et al. Ann. N.Y. Acad. Sci. 1291, 14-32 (2013).
- 6. Yunokawa, M., Koizumi, F., Kitamura, Y., et al. Cancer Sci. 103(9), 1665-1671 (2012).
- 7. Gurk-Turner, C., Manitpisitkul, W., and Cooper, M. Transplantation 94(7), 659-668 (2012).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information Buyer agrees to purchase the material can be found on our website.

Copyright Cayman Chemical Company, 11/04/2022

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM