



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



PRODUCT INFORMATION

L- α -Hydroxyglutaric Acid (sodium salt)

Item No. 11876

CAS Registry No.: 63512-50-5

Formal Name: 2S-hydroxy-pantanedioic acid, disodium salt

Synonyms: L-2-HG, L-2-Hydroxyglutarate, L-2-Hydroxyglutaric Acid, (S)-2-Hydroxypantanedioic Acid

MF: C₅H₈O₅ • 2Na


FW: 194.1

Purity: ≥95%

Supplied as: A crystalline solid

Storage: -20°C

Stability: ≥2 years



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

L- α -Hydroxyglutaric acid (sodium salt) is supplied as a crystalline solid. Aqueous solutions of L- α -hydroxyglutaric acid (sodium salt) can be prepared by directly dissolving the crystalline solid in aqueous buffers. The solubility of L- α -hydroxyglutaric acid (sodium salt) in PBS, pH 7.2, is approximately 10 mg/ml. We do not recommend storing the aqueous solution for more than one day.

Description

L- α -Hydroxyglutaric acid (L-2-HG) is an α -hydroxy acid. It is metabolized to 2-oxoglutarate (α -ketoglutarate) by L-2-hydroxyglutarate dehydrogenase, and mutations in this enzyme lead to 2-hydroxyglutaric aciduria, a neurometabolic disorder characterized by increased L-2-HG levels.^{1,2} L-2-HG is structurally similar to α -ketoglutarate and competitively inhibits α -ketoglutarate-dependent dioxygenases, including several involved in histone lysine and DNA demethylation.^{3,4}

References

- Rzem, R., Veiga-da-Cunha, M., Noël, G., et al. A gene encoding a putative FAD-dependent L-2-hydroxyglutarate dehydrogenase is mutated in L-2-hydroxyglutaric aciduria. *Proc. Nat. Acad. Sci. USA* **101**(48), 16849-16854 (2004).
- Struys, E.A., Verhoeven, N.M., Roos, B., et al. Disease-related metabolites in culture medium of fibroblasts from patients with D-2-hydroxyglutaric aciduria, L-2-hydroxyglutaric aciduria, and combined D/L-2-hydroxyglutaric aciduria. *Clin. Chem.* **49**(7), 1133-1138 (2003).
- Chowdhury, R., Yeoh, K.K., Tian, Y.M., et al. The oncometabolite 2-hydroxyglutarate inhibits histone lysine demethylases. *EMBO Rep.* **12**(5), 463-469 (2011).
- Xu, W., Yang, H., Liu, Y., et al. Oncometabolite 2-hydroxyglutarate is a competitive inhibitor of α -ketoglutarate-dependent dioxygenases. *Cancer Cell.* **19**(1), 17-30 (2011).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 10/26/2018

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 - USA

PHONE: [800] 364-9897
[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM