



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 


PRODUCT INFORMATION



15-deoxy- $\Delta^{12,14}$ -Prostaglandin D₂

Item No. 12700

CAS Registry No.: 85235-11-6
Formal Name: 9 α -hydroxy-11-oxo-prosta-5Z,12E,14E-trien-1-oic acid
Synonym: 15-deoxy- $\Delta^{12,14}$ -PGD₂
MF: C₂₀H₃₀O₄
FW: 334.5
Purity: \geq 95%
UV/Vis.: λ_{\max} : 296 nm
Supplied as: A solution in methyl acetate
Storage: -20°C
Stability: \geq 1 year



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

15-deoxy- $\Delta^{12,14}$ PGD₂ is supplied as a solution in methyl acetate. To change the solvent, simply evaporate the methyl acetate under a gentle stream of nitrogen and immediately add the solvent of choice. Solvents such as DMSO, dimethyl formamide, or ethanol purged with an inert gas can be used. The solubility of 15-deoxy- $\Delta^{12,14}$ PGD₂ in these solvents is approximately 50 mg/ml.

Further dilutions of the stock solution into aqueous buffers or isotonic saline should be made prior to performing biological experiments. Ensure that the residual amount of organic solvent is insignificant, since organic solvents may have physiological effects at low concentrations. If an organic solvent-free solution of 15-deoxy- $\Delta^{12,14}$ PGD₂ is needed, it can be prepared by evaporating the methyl acetate and directly dissolving the neat oil in aqueous buffers. The solubility of 15-deoxy- $\Delta^{12,14}$ PGD₂ in PBS, pH 7.2, is approximately 5 mg/ml. We do not recommend storing the aqueous solution for more than one day.

15-deoxy- $\Delta^{12,14}$ PGD₂ is a synthetic analog of PGD₂ and a potential precursor to the PPAR γ ligand 15-deoxy- $\Delta^{12,14}$ PGJ₂.³ 15-deoxy- $\Delta^{12,14}$ PGD₂ possesses cytotoxic effects on L1210 mouse leukemia cells *in vitro* with an IC₅₀ of 0.3 μ g/ml. 15-deoxy- $\Delta^{12,14}$ PGD₂ is ten-fold more potent than PGD₂ in this activity.¹ However, 15-deoxy- $\Delta^{12,14}$ PGD₂ is a less effective inhibitor of ADP-induced platelet aggregation (IC₅₀ = 320 ng/ml) compared to PGD₂ (IC₅₀ = 3 ng/ml).²

Description

15-deoxy- $\Delta^{12,14}$ -PGD₂ is a metabolite of PGD₂ (Item No. 12010).¹ It is an agonist of PGD₂ receptor 2 (DP₂) that binds DP₂ (K_i = 50 nM for the mouse receptor expressed in HEK293 cell membranes) and induces activation of eosinophils (EC₅₀ = 8 nM).^{2,3} It also stimulates the recruitment of steroid receptor coactivator-1 (SRC-1) to peroxisome proliferator-activated receptor γ (PPAR γ) and induces PPAR γ -mediated transcription in a reporter assay when used at a concentration of 5 μ M.¹ 15-deoxy- $\Delta^{12,14}$ -PGD₂ is cytotoxic to L1210 murine leukemia cells (IC₅₀ = 0.3 μ g/ml).⁴ It inhibits ADP-induced platelet aggregation (IC₅₀ = 320 ng/ml) less potently than PGD₂.⁵

References

1. Söderström, M., Wigren, J., Surapureddi, S., *et al.* *Biochim. Biophys. Acta* **1631**(1), 35-41 (2003).
2. Hata, A.N., Zent, R., Breyer, M.D., *et al.* *J. Pharmacol. Exp. Ther.* **306**(2), 463-470 (2003).
3. Monneret, G., Li, H., Vasilescu, J., *et al.* *J. Immunol.* **168**(7), 3563-3569 (2002).
4. Forman, B.M., Tontonoz, P., Chen, J., *et al.* *Cell* **83**(5), 803-812 (1995).
5. Bundy, G.L., Morton, D.R., Peterson, D.C., *et al.* *J. Med. Chem.* **26**(6), 790-799 (1983).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 09/10/2020

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD

ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM

WWW.CAYMANCHEM.COM