



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com


[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

PRODUCT INFORMATION



Prostaglandin E₂ Item No. 14010

CAS Registry No.: 363-24-6
Formal Name: 9-oxo-11 α ,15S-dihydroxy-prosta-5Z,13E-dien-1-oic acid
Synonyms: Dinoprostone, PGE₂
MF: C₂₀H₃₂O₅
FW: 352.5
Purity: \geq 98%
Supplied as: A crystalline solid
Storage: -20°C
Stability: \geq 2 years



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

Prostaglandin E₂ (PGE₂) is supplied as a crystalline solid. A stock solution may be made by dissolving the PGE₂ in the solvent of choice. PGE₂ is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide, which should be purged with an inert gas. The solubility of PGE₂ in these solvents is >100 mg/ml.

Further dilutions of the stock solution into aqueous buffers or isotonic saline should be made prior to performing biological experiments. Ensure that the residual amount of organic solvent is insignificant, since organic solvents may have physiological effects at low concentrations. Organic solvent-free aqueous solutions of PGE₂ can be prepared by directly dissolving the crystalline solid in aqueous buffers. The solubility of PGE₂ in PBS, pH 7.2, is approximately 5 mg/ml. We do not recommend storing the aqueous solution for more than one day.

Description

PGE₂ is one of the primary cyclooxygenase products of arachidonic acid (Item No. 90010) and one of the most widely investigated PGs. Its activity influences inflammation, fertility and parturition, gastric mucosal integrity, and immune modulation.¹⁻⁴ The effects of PGE₂ are transduced by at least four distinct receptors designated EP₁, EP₂, EP₃, and EP₄.⁵ Affinity constants (K_d) of PGE₂ for these receptors range from 1-10 nM depending on the receptor subtype and tissue.

References

1. Willis, A.L. and Cornelsen, M. Repeated injection of prostaglandin E₂ in rat paws induces chronic swelling and a marked decrease in pain threshold. *Prostaglandins* **3(3)**, 353-357 (1973).
2. Jackson, G.M., Sharp, H.T., and Varner, M.W. Cervical ripening before induction of labor: A randomized trial of prostaglandin E₂ gel versus low-dose oxytocin. *Am. J. Obstet. Gynecol.* **171(4)**, 1092-1096 (1994).
3. Robert, A., Schultz, J.R., Nezamis, J.E., et al. Gastric antisecretory and antiulcer properties of PGE₂, 15-methyl PGE₂, and 16,16-dimethyl PGE₂. Intravenous, oral and intrajejunal administration. *Gastroenterology* **70(3)**, 359-370 (1976).
4. Arvind, P., Papavassiliou, E.D., Tsioulas, G.J., et al. Prostaglandin E₂ down-regulates the expression of HLA-DR antigen in human colon adenocarcinoma cell lines. *Biochemistry* **34(16)**, 5604-5609 (1995).
5. Coleman, R.A., Smith, W.L., and Narumiya, S. Classification of prostanoid receptors: Properties, distribution, and structure of the receptors and their subtypes. *Pharmacol. Rev.* **46(2)**, 205-229 (1994).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 03/15/2018

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM