



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 


PRODUCT INFORMATION



13,14-dihydro-15-keto Prostaglandin E₂

Item No. 14650

CAS Registry No.: 363-23-5
Formal Name: (5Z,11α)-11-hydroxy-9,15-dioxo-prost-5-en-1-oic acid
Synonyms: 13,14-dihydro-15-keto PGE₂, 13,14-dihydro-oxo-PGE₂, PGEM
MF: C₂₀H₃₂O₅
FW: 352.5
Purity: ≥95%
Supplied as: A solution in methyl acetate
Storage: -20°C
Stability: ≥2 years



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

13,14-dihydro-15-keto Prostaglandin E₂ (13,14-dihydro-15-keto PGE₂) is supplied as a solution in methyl acetate. To change the solvent, simply evaporate the methyl acetate under a gentle stream of nitrogen and immediately add the solvent of choice. Solvents such as ethanol, DMSO, and dimethyl formamide purged with an inert gas can be used. The solubility of 13,14-dihydro-15-keto PGE₂ in these solvents is approximately 100 mg/ml.

Further dilutions of the stock solution into aqueous buffers or isotonic saline should be made prior to performing biological experiments. Ensure that the residual amount of organic solvent is insignificant, since organic solvents may have physiological effects at low concentrations. If an organic solvent-free solution of 13,14-dihydro-15-keto PGE₂ is needed, it can be prepared by evaporating the methyl acetate and directly dissolving the neat oil in aqueous buffers. The solubility of 13,14-dihydro-15-keto PGE₂ in PBS (pH 7.2) is approximately 5 mg/ml. We do not recommend storing the aqueous solution for more than one day.

Description

13,14-dihydro-15-keto PGE₂ is a metabolite of PGE₂ (Item No. 14010) and the primary PGE₂ metabolite in plasma.^{1,2} It is formed from PGE₂ via a 15-keto PGE₂ intermediate by 15-oxo-PG Δ¹³ reductase.² Unlike PGE₂, 13,14-dihydro-15-keto PGE₂ does not bind effectively to the PGE₂ receptors EP₂ and EP₄ expressed in CHO cells (K_s = 12 and 57 μM, respectively) or induce adenylate cyclase activity in the same cells (EC₅₀s = >18 and >38 μM, respectively). Levels of 13,14-dihydro-15-keto PGE₂ are increased in the plasma of women in the third trimester of pregnancy and in women during and immediately after labor and delivery.³ Levels of 13,14-dihydro-15-keto PGE₂ levels are decreased in tumor tissue compared to adjacent non-cancerous tissue isolated from patients with non-small cell lung cancer (NSCLC).⁴

References

1. Hamberg, M. and Samuelsson, B. On the metabolism of prostaglandins E₁ and E₂ in man. *J. Biol. Chem.* **246**(22), 6713-6721 (1971).
2. Nishigaki, N., Negishi, M., and Ichikawa, A. Two G_s-coupled prostaglandin E receptor subtypes, EP2 and EP4, differ in desensitization and sensitivity to the metabolic inactivation of the agonist. *Mol. Pharmacol.* **50**(4), 1031-1037 (1996).
3. Husslein, P. and Sinzinger, H. Concentration of 13,14-dihydro-15-keto-prostaglandin E₂ in the maternal peripheral plasma during labour of spontaneous onset. *Br. J. Obstet. Gynaecol.* **91**(3), 228-231 (1984).
4. Hughes, D., Otani, T., Yang, P., et al. NAD⁺-dependent 15-hydroxyprostaglandin dehydrogenase regulates levels of bioactive lipids in non-small cell lung cancer. *Cancer Prev. Res. (Phila)* **1**(4), 241-249 (2008).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 11/30/2021

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897
[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM