

## Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

#### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com

# **PRODUCT** INFORMATION



#### Exemestane

Item No. 15008

CAS Registry No.:	107868-30-4	
Formal Name:	6-methylene-androsta-1,4-diene-	• 0
	3,17-dione	$\sim \mathcal{I}$
Synonyms:	Aromasin™, FCE 24304	$\int \int \langle \nabla $
MF:	$C_{20}H_{24}O_{2}$	$\land$ $\blacksquare$ $\downarrow$ $\downarrow$ $\downarrow$ $\downarrow$ /
FW:	296.4	
Purity:	≥95%	
UV/Vis.:	λ <sub>max</sub> : 246 nm	0
Supplied as:	A crystalline solid	CH-
Storage:	-20°C	012
Stability:	≥2 years	

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

#### Laboratory Procedures

Exemestane is supplied as a crystalline solid. A stock solution may be made by dissolving the exemestane in the solvent of choice. Exemestane is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide (DMF), which should be purged with an inert gas. The solubility of exemestane in ethanol is approximately 20 mg/ml and approximately 30 mg/ml in DMSO and DMF.

Exemestane is sparingly soluble in aqueous buffers. For maximum solubility in aqueous buffers, exemestane should first be dissolved in DMSO and then diluted with the aqueous buffer of choice. Exemestane has a solubility of approximately 0.5 mg/ml in a 1:1 solution of DMSO:PBS (pH 7.2) using this method. We do not recommend storing the aqueous solution for more than one day.

#### Description

Exemestane is a third generation, irreversible steroidal aromatase inhibitor (K<sub>i</sub> = 10.2 nM;  $K_{inact}$  = 26 nM) that induces aromatase degradation leading to a decrease in estrogen levels in plasma.<sup>1,2</sup> As an androgen analog, exemestane exhibits androgenic effects and has been shown to decrease total and HDL cholesterol, apo A₁, and total triglyceride levels.<sup>3</sup> Marketed under the trade name Aromasin<sup>™</sup>, exemestane has been used to treat estrogen receptor-positive breast cancers in post-menopausal women.<sup>1</sup>

#### References

- 1. Wang, X. and Chen, S. Aromatase destabilizer: Novel action of exemestane, a food and drug administration-approved aromatase inhibitor. Cancer Res. 66(21), 10281-10286 (2006).
- 2. Santen, R.J. and Harvey, H.A. Use of aromatase inhibitors in breast carcinoma. Endocr. Relat. Cancer 6(1), 75-92 (1999).
- 3. Buzdar, A.U., Robertson, J.F.R., Eiermann, W., et al. An overview of the pharmacology and pharmacokinetics of the newer generation aromatase inhibitors anastrozole, letrozole, and exemestane. Cancer 95(9), 2006-2016 (2002).

WARNING THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

#### SAFFTY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

#### WARRANTY AND LIMITATION OF REMEDY

uyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 11/05/2018

#### CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897 [734] 971-3335 FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.CAYMANCHEM.COM