

Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com

Product Information



1,2-Distearoyl-sn-glycerol

Item No. 15079

CAS Registry No.:	10567-21-2	
Formal Name:	1,2-distearoyl-sn-glycerol	0
Synonyms:	1,2-Dioctadecanoyl-sn-glycerol,	
	DSG	о О
MF:	C ₃₉ H ₇₆ O ₅	$ \land \land$
FW:	625.0	
Purity:	≥98%	
Stability:	≥2 years at -20°C	он
Supplied as:	A crystalline solid	

Laboratory Procedures

For long term storage, we suggest that 1,2-distearoyl-sn-glycerol (DSG) be stored as supplied at -20°C. It should be stable for at least two years.

DSG is supplied as a crystalline solid. A stock solution may be made by dissolving the DSG in the solvent of choice. DSG is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide, which should be purged with an inert gas. The solubility of DSG in these solvents is approximately 0.25, 30, and 20 mg/ml, respectively.

Further dilutions of the stock solution into aqueous buffers or isotonic saline should be made prior to performing biological experiments. Ensure that the residual amount of organic solvent is insignificant, since organic solvents may have physiological effects at low concentrations. Organic solvent-free aqueous solutions of DSG can be prepared by directly dissolving the crystalline solid in aqueous buffers. The solubility of DSG in PBS, pH 7.2, is approximately 0.7 mg/ml. We do not recommend storing the aqueous solution for more than one day.

1,2-Diacylglycerols (DAGs) are formed in cells through the hydrolysis of phospatidyliositol 4,5-bisphosphate by phospholipase C. They have important roles in signal transduction, as in their activation of some isoforms of PKC.¹ In addition, DAGs are modified by DAG kinases, reducing signaling through DAG while generating bioactive phosphatidic acids.^{2,3} DSG is a form of DAG containing the saturated long-chain (18:0) stearic acid at both the sn-1 and the sn-2 position.

References

- 1. Nakamura, S. and Nishizuka, Y. Lipid mediators and protein kinase C activation for the intracellular signaling network. J. Biochem. 115(6), 1029-1034 (1994).
- 2. Topham, M.K. and Prescott, S.M. Mammalian diacylglycerol kinases, a family of lipid kinases with signaling functions. J. Biol. Chem. 274, 11447-11450 (1999).
- 3. Joshi, R.P. and Koretzky, G.A. Diacylglycerol kinases: Regulated controllers of T cell activation, function, and development. Int. J. Mol. Sci. 14(4), 6649-6673 (2013).

Related Products

For a list of related products please visit: www.caymanchem.com/catalog/15079

WARNING: This product is for laboratory research only: not for administration to humans. Not for human or veterinary DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until information to the contrary becomes available. Do not ingest, swallow, or inhale. Do not get in eyes, on skin, or on clothing. Wash thoroughly after handling. This information contains some, but not all, of the information required for the safe and proper use of this material. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Cayman Chemical Company makes no warranty or guarantee of any kind, whether written or oral, expressed or implied, including without limitation, any warranty of fitness for a particular purpose, suitability and merchantability, which extends beyond the description of the chemicals hereof. Cayman warrants only to the original customer that the material will meet our specifications at the time of delivery.

ar the time of delivery. Cayman will carry out its delivery obligations with due care and skill. Thus, in no event will Cayman have **any obligation or liability**, whether in tort (including negligence) or in contract, for any direct, indirect, indirect incidental or consequential damages, even if Cayman is informed about their possible existence. This limitation of liability does not apply in the case of intentional acts or negligence of Cayman, its directors or its employees. Buyer's **exclusive remedy** and Cayman's sole liability hereunder shall be limited to a <u>refund</u> of the purchase price, or at Cayman's option, the <u>replacement</u>, at no cost to Buyer, of all material that

Buyer's exclusive remedy and Laymans sole hability neterinder shall be infinited to a terminal of the particular process of an experimental strength of the particular process of the particular proces of the

Phone (800) 364-9897

Cayman Chemical

Mailing address 1180 E. Ellsworth Road Ann Arbor, MI 48108 USA

(734) 971-3335 Fax

(734) 971-3640

E-Mail

custserv@caymanchem.com

Web

www.cavmanchem.com