



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 


PRODUCT INFORMATION



1,2-Dipalmitoyl-*sn*-glycero-O-ethyl-3-PC (chloride)

Item No. 15099

CAS Registry No.: 328250-18-6
Formal Name: 4-ethoxy-N,N,N-trimethyl-10-oxo-7R-[(1-oxohexadecyl)oxy]-4-oxide-3,5,9-trioxa-4-phosphapentacosan-1-aminium, monochloride
Synonyms: DPePC, 1,2-Dipalmitoyl-*sn*-glycero-O-ethyl-3-Phosphocholine, 1,2-EDPPC
MF: C₄₂H₈₅NO₈P • Cl
FW: 798.6
Purity: ≥95%
Supplied as: A crystalline solid
Storage: -20°C
Stability: ≥2 years



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

1,2-Dipalmitoyl-*sn*-glycero-O-ethyl-3-PC (chloride) is supplied as a crystalline solid. A stock solution may be made by dissolving the 1,2-dipalmitoyl-*sn*-glycero-O-ethyl-3-PC (chloride) in the solvent of choice, which should be purged with an inert gas. 1,2-Dipalmitoyl-*sn*-glycero-O-ethyl-3-PC (chloride) is soluble in organic solvents such as ethanol and chloroform. The solubility of 1,2-dipalmitoyl-*sn*-glycero-O-ethyl-3-PC (chloride) in these solvents is approximately 25 and 10 mg/ml, respectively.

Description

1,2-Dipalmitoyl-*sn*-glycero-O-ethyl-3-PC is a phospholipid containing the saturated long-chain (16:0) stearic acid inserted at the *sn*-1 and *sn*-2 positions and an alkyl group on the phosphate oxygen of the polar headgroup. Cationic phospholipids such as this have proved useful as DNA transfection agents and for studies of surface charge density within lipid bilayers.^{1,2}

References

1. Lewis, R.N., Winter, I., Kriechbaum, M., *et al.* Studies of the structure and organization of cationic lipid bilayer membranes: Calorimetric, spectroscopic, and x-ray diffraction studies of linear saturated P-O-ethyl phosphatidylcholines. *Biophys. J.* **80**(3), 1329-1342 (2001).
2. Kopin, I.J., Axelrod, J., and Gordon, E. The metabolic fate of H³-epinephrine and C¹⁴-metanephrine in the rat. *J. Biol. Chem.* **236**(7), 2109-2136 (1961).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 10/12/2021

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897
[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM