



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# PRODUCT INFORMATION



## 6,15-diketo-13,14-dihydro Prostaglandin F<sub>1α</sub>

Item No. 15270

CAS Registry No.: 63983-53-9

Formal Name: 6,15-dioxo-9α,11α-dihydroxy-prostan-1-oic acid

Synonym: 6,15-diketo-13,14-dihydro PGF<sub>1α</sub>

MF: C<sub>20</sub>H<sub>34</sub>O<sub>6</sub>


FW: 370.5

Purity: ≥98% (mixture of tautomers)

Supplied as: A solution in methyl acetate

Storage: -20°C

Stability: As supplied, 1 year from the QC date provided on the Certificate of Analysis, when stored properly



### Laboratory Procedures

6,15-diketo-13,14-dihydro PGF<sub>1α</sub> is supplied as a solution in methyl acetate. To change the solvent, simply evaporate the methyl acetate under a gentle stream of nitrogen and immediately add the solvent of choice. Solvents such as ethanol, DMSO, or dimethyl formamide purged with an inert gas can be used. The solubility of 6,15-diketo-13,14-dihydro PGF<sub>1α</sub> in these solvents is approximately 16 mg/ml. 6,15-diketo-13,14-dihydro PGF<sub>1α</sub> is stable for at least six months in these solvents if stored at -20°C.

Further dilutions of the stock solution into aqueous buffers or isotonic saline should be made prior to performing biological experiments. Ensure that the residual amount of organic solvent is insignificant, since organic solvents may have physiological effects at low concentrations. If an organic solvent-free aqueous solution of 6,15-diketo-13,14-dihydro PGF<sub>1α</sub> is needed, it can be prepared by evaporating the methyl acetate and directly dissolving the neat oil in aqueous buffers. The solubility of 6,15-diketo-13,14-dihydro PGF<sub>1α</sub> in PBS (pH 7.2) is approximately 3.3 mg/ml. Store aqueous solutions of 6,15-diketo-13,14-dihydro PGF<sub>1α</sub> on ice and use within 12 hours of preparation.

### Description

6,15-diketo-13,14-dihydro PGF<sub>1α</sub> is a metabolite of PGI<sub>2</sub>. It was shown to enhance intracellular cAMP and cholesterol catabolism in bovine arterial smooth muscle cells.<sup>1</sup>

### Reference

1. Etingin, O.R., Weksler, B.B., and Hajjar, D.P. Cholesterol metabolism is altered by hydrolytic metabolites of prostacyclin in arterial smooth muscle cells. *J. Lipid Res.* **27**, 530-536 (1986).

#### WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

#### SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

#### WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 03/21/2016

#### CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD  
ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM

WWW.CAYMANCHEM.COM