

Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com

PRODUCT INFORMATION



Kasugamycin (hydrochloride)

Item No. 15322

CAS Desister No. 10400 44 0

CAS Registry No.:	19408-46-9	
Formal Name:	3-0-[2-amino-4-	
	[(carboxyiminomethyl)amino]-	QH NH ₂
	2,3,4,6-tetradeoxy-α-D-arabino-	HOOO
	hexopyranosyl]-D-chiro-inositol,	Υ Υ Υ Υ
	monohydrochloride	NH II
MF:	$C_{14}H_{25}N_3O_9 \bullet HCI$	HO OH OH OH
FW:	415.8	
Purity:	≥70%	он н о
Supplied as:	A crystalline solid	
Storage:	-20°C	
Stability:	≥4 years	
Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.		

Laboratory Procedures

Kasugamycin (hydrochloride) is supplied as a crystalline solid. Aqueous solutions of kasugamycin (hydrochloride) can be prepared by directly dissolving the crystalline solid in aqueous buffers. The solubility of kasugamycin (hydrochloride) in PBS, pH 7.2, is approximately 5 mg/ml. We do not recommend storing the aqueous solution for more than one day.

Description

Kasugamycin is an aminoglycoside that has been found in S. kasugaensis and has diverse biological activities.¹⁻³ It is active against S. dysenteriae, ¹ Kasugamycin (1 kg/ha) reduces the severity of rice sheath blight induced by the plant pathogenic fungi R. solani and C. janseana-induced rice narrow brown leaf spot, as well as increases rice yield.² It also inhibits acidic mammalian chitinase (CHIA), also known as AMCase, with an IC₅₀ value of 4.13 μ M. Kasugamycin (12.5, 25, 50, or 100 mg/kg every other day) decreases lung collagen levels in a mouse model of pulmonary fibrosis induced by the glycopeptide antitumor antibiotic bleomycin (Item No. 13877).³ Formulations containing kasugamycin have been used as fungicides in agriculture.

References

- 1. Hamada, M., Hashimoto, T., Takahashi, T., et al. Antimicrobial activity of kasugamycin. J. Antibiot. (Tokyo) 18, 104-106 (1965).
- 2. Uppala, S. and Zhou, X.G. Field efficacy of fungicides for management of sheath blight and narrow brown leaf spot of rice. Crop Prot. 104, 72-77 (2018).
- 3. Lee, J.-H., Lee, C.-M., Lee, J.H., et al. Kasugamycin is a novel chitinase 1 inhibitor with strong antifibrotic effects on pulmonary fibrosis. Am. J. Respir. Cell Mol. Biol. 67(3), 309-319 (2022).

WARNING THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFFTY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 7/28/2023

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897 [734] 971-3335 FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.CAYMANCHEM.COM