

Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com

PRODUCT INFORMATION



Validamycin A

Item No. 15326

CAS Registry No.:	37248-47-8		
Formal Name:	1,5,6-trideoxy-4-O-β-D-glucopyranosyl-5-(hydroxymethyl)-	но	HO
	1-[[(1S,4R,5S,6S)-4,5,6-trihydroxy-3-(hydroxymethyl)-2- cyclobeyen-1-yllaminol-D-chiro-inositol		ОН
Curra and read	Velidesin	HU.	
Synonym:	Validacin		
MF:	C ₂₀ H ₃₅ NO ₁₃		
FW:	497.5	но	N OH O
Purity:	≥60%		
Supplied as:	A crystalline solid	Оп	нон
Storage:	-20°C		`OH
Stability:	≥4 years		
Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis			

Laboratory Procedures

Validamycin A is supplied as a crystalline solid. A stock solution may be made by dissolving the validamycin A in the solvent of choice, which should be purged with an inert gas. Validamycin A is soluble in the organic solvent DMSO at a concentration of approximately 2 mg/ml.

Further dilutions of the stock solution into aqueous buffers or isotonic saline should be made prior to performing biological experiments. Ensure that the residual amount of organic solvent is insignificant, since organic solvents may have physiological effects at low concentrations. Organic solvent-free aqueous solutions of validamycin A can be prepared by directly dissolving the crystalline solid in aqueous buffers. The solubility of validamycin A in PBS (pH 7.2) is approximately 2 mg/ml. We do not recommend storing the aqueous solution for more than one day.

Description

Validamycin A is a fungicidal agent used to protect rice plants against sheath blight caused by the pathogenic fungus, R. solani, due to its ability to inhibit trehalase, a trehalose-hydrolizing enzyme ($K_i = 1.9 \text{ nM}$; IC₅₀ = 0.7 μ M).¹ It is also effective against the growth and sporulation of *R. cerealis*, *F. culmorum*, and other fungi.² This compound has also been used to study trehalase activity and trehalose biosynthesis.³

References

- 1. Asano, N., Yamaguchi, T., Kameda, Y., et al. Effect of validamycins on glycohydrolases of Rhizoctonia solani. J. Antibiot. (Tokyo) 40(4), 526-532 (1987).
- 2. Robson, G.D., Kuhn, P.J., and Trinci, A.P. Effects of validamycin A on the morphology, growth and sporulation of Rhizoctonia cerealis, Fusarium culmorum and other fungi. J. Gen. Microbiol. 134(12), 3187-3194 (1988).
- 3. Goddijn, O.J., Verwoerd, T.C., Voogd, E., et al. Inhibition of trehalase activity enhances trehalose accumulation in transgenic plants. Plant Physiol. 113(1), 181-190 (1997).

WARNING THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

uyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 07/26/2023

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897 [734] 971-3335 FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.CAYMANCHEM.COM