

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



3-Hydroxyglutaric Acid

Item No. 16334

CAS Registry No.: 638-18-6

Formal Name: 3-hydroxy-pentanedioic acid Synonyms: 2,4-Dideoxypentaric Acid, 3-HG,

3-Hydroxyglutarate, β-Hydroxyglutaric Acid,

3-Hydroxypentanedioic Acid, 3-OH-GA

MF: $C_5H_8O_5$ FW: 148.1 **Purity:** ≥95%

Stability: ≥2 years at -20°C Supplied as: A crystalline solid

Laboratory Procedures

For long term storage, we suggest that 3-hydroxyglutaric acid be stored as supplied at -20°C. It should be stable for at least two years.

3-Hydroxyglutaric acid is supplied as a crystalline solid. A stock solution may be made by dissolving the 3-hydroxyglutaric acid in the solvent of choice. 3-Hydroxyglutaric acid is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide, which should be purged with an inert gas. The solubility of 3-hydroxyglutaric acid in these solvents is approximately 30 mg/ml.

Further dilutions of the stock solution into aqueous buffers or isotonic saline should be made prior to performing biological experiments. Ensure that the residual amount of organic solvent is insignificant, since organic solvents may have physiological effects at low concentrations. Organic solvent-free aqueous solutions of 3-hydroxyglutaric acid can be prepared by directly dissolving the crystalline solid in aqueous buffers. The solubility of 3-hydroxyglutaric acid in PBS, pH 7.2, is approximately 10 mg/ml. We do not recommend storing the aqueous solution for more than one day.

Description

Glutaryl-CoA dehydrogenase (GCDH) is a mitochondrial matrix protein that catalyzes the oxidative decarboxylation of glutaryl-CoA to crotonyl-CoA and carbon dioxide in the catabolic pathways of lysine. hydroxylysine, and tryptophan metabolism. 3-Hydroxyglutaric acid is one of several metabolites produced when insufficient levels of GCDH are available. 1,2 Urinary levels of 3-hydroxyglutaric acid are elevated during glutaric acidemia type 1, an autosomal recessive GCDH deficiency disorder that can lead to neurodegeneration if left untreated. 1,2 3-Hydroxyglutaric acid is used as a biomarker of GCDH deficiency. 1,2

References

- 1. Fang, N. and Casida, J.E. Anticancer action of cubé insecticide: Correlation for rotenoid constituents between inhibition of NADH: Ubiquinone oxidoreductase and induced ornithine decarboxylase activities. Proc. Natl. Acad. Sci. USA 95, 3380-3384 (1998).
- 2. Sauer, S.W., Okun, J.G., Schwab, M.A., et al. Bioenergetics in glutaryl-coenzyme a dehydrogenase deficiency. A role for glutaryl-coenzyme A. J. Biol. Chem. 290(23), 21830-21836 (2005).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

al should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 09/22/2015

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA **PHONE:** [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM