

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



2,4-Diacetylphloroglucinol

Item No. 16345

CAS Registry No.: 2161-86-6

Formal Name: 1,1'-(2,4,6-trihydroxy-1,3-phenylene)bis-ethanone Synonyms: 5-Acetyl-2,4,6-trihydroxyacetophenone, DAPG,


1,3-Diacetyl-2,4,6-trihydroxybenzene

MF: $C_{10}H_{10}O_5$ 210.2 FW: ≥98% **Purity:**

 λ_{max} : 269, 329 nm UV/Vis.: Supplied as: A crystalline solid

Storage: -20°C Stability: ≥4 years

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.



Laboratory Procedures

2,4-Diacetylphloroglucinol (DAPG) is supplied as a crystalline solid. A stock solution may be made by dissolving the DAPG in the solvent of choice, which should be purged with an inert gas. DAPG is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide (DMF). The solubility of DAPG in these solvents is approximately 25, 16, and 30 mg/ml, respectively.

DAPG is sparingly soluble in aqueous buffers. For maximum solubility in aqueous buffers, DAPG should first be dissolved in DMF and then diluted with the aqueous buffer of choice. DAPG has a solubility of approximately 0.5 mg/ml in a 1:1 solution of DMF:PBS (pH 7.2) using this method. We do not recommend storing the aqueous solution for more than one day.

Description

2,4-Diacetylphloroglucinol is a bacterial metabolite originally isolated from P. fluorescens that has diverse biological activities. It is active against B. subtillis, E. faecalis, S. aureus, and S. lividans (MICs = 5, 50, 5, and 5 µg/ml, respectively). 2,4-Diacetylphloroglucinol is also active against the plant pathogenic fungi F. oxysporum, P. debaryanum, and R. solani (MICs = 16, 64, and 32-64 µg/ml, respectively). It reduces the growth of corn, flax, tobacco, and tomato seedlings (MICs = 16-32, 32, 8, and 16 µg/ml, respectively) and suppresses T. basicola-induced black root rot on tobacco plants. It also inhibits cell growth in A549 lung, MDA-MB-231 breast, and HeLa cervical cancer cells (IC₅₀s = 52, 94, and 141 nM, respectively).²

References

- 1. Keel, C., Ursula, S., Maurhofer, M., et al. Suppression of root diseases by Pseudomonas fluorescens CHAO: Importance of the bacterial seconday metabolite 2,4-diacetylphloroglucinol. Mol. Plant Microbe Interact. **5(1)**, 4-13 (1995).
- 2. Veena, V.K., Popavath, R.N., Kennedy, K., et al. In vitro antiproliferative, pro-apoptotic, antimetastatic and anti-inflammatory potential of 2,4-diacetylphloroglucinol (DAPG) by Pseudomonas aeruginosa strain FP10. Apoptosis 20(10), 1281-1295 (2015).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information Buyer agrees to purchase the mater can be found on our website.

Copyright Cayman Chemical Company, 02/02/2023

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM