

## Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

#### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com

# **PRODUCT** INFORMATION



**19-alkyne Cholesterol** 

Item No. 16446

| CAS Registry No.:<br>Formal Name: | 2058319-71-2<br>10-ethynyl-19-norcholest-5-en-3β-ol | ``,   |
|-----------------------------------|---|-------|
| Synonym:                          | 19-ethynyl Cholesterol                              | A Lot |
| MF:                               | C <sub>28</sub> H <sub>44</sub> O                   |       |
| FW:                               | 396.7   | н /   |
| Purity:                           | ≥95%  |       |
| Supplied as:                      | A crystalline solid                                 | н н   |
| Storage:                          | -20°C   |       |
| Stability:                        | ≥2 years  | HO    |
|                                   |   |       |

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

#### Laboratory Procedures

19-alkyne Cholesterol is supplied as a crystalline solid. A stock solution may be made by dissolving the 19-alkyne cholesterol in the solvent of choice, which should be purged with an inert gas. 19-alkyne Cholesterol is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide. The solubility of 19-alkyne cholesterol in these solvents is approximately 20, 0.1, and 2 mg/ml, respectively.

19-alkyne Cholesterol is sparingly soluble in aqueous buffers. For maximum solubility in aqueous buffers, 19-alkyne Cholesterol should first be dissolved in ethanol and then diluted with the aqueous buffer of choice. 19-alkyne Cholesterol has a solubility of approximately 0.3 mg/ml in a 1:2 solution of ethanol:PBS (pH 7.2) using this method. We do not recommend storing the aqueous solution for more than one day.

#### Description

19-alkyne Cholesterol is an alkyne derivative of cholesterol (Item No. 9003100).<sup>1</sup> It has been used to track cholesterol localization and subcellular distribution in NIH3T3 cells and to study the role of cholesterol in hedgehog (Hh) signal transduction in Shh-LIGHT2 cells using click chemistry and various reporter azides.

#### Reference

1. Jao, C.Y., Nedelcu, D., Lopez, L.V., et al. Bioorthogonal probes for imaging sterols in cells. Chembiochem. 16(4), 611-617 (2015).

WARNING THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

#### SAFFTY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

#### WARRANTY AND LIMITATION OF REMEDY

uyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 02/07/2022

#### CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897 [734] 971-3335 FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.CAYMANCHEM.COM