

Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com


Product informat



Erythromycin

Item No. 16486

CAS Registry No.:	114-07-8
Formal Name:	erythromycin
Synonym:	NSC 55929
MF:	C ₃₇ H ₆₇ NO ₁₃
FW:	733.9
Purity:	≥98%
Supplied as:	A crystalline solid
Storage:	-20°C
Stability:	≥2 years



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

Erythromycin is supplied as a crystalline solid. A stock solution may be made by dissolving the erythromycin in the solvent of choice. Erythromycin is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide (DMF), which should be purged with an inert gas. The solubility of erythromycin is approximately 30 mg/ml in ethanol and 15 mg/ml in DMSO and DMF.

Erythromycin is sparingly soluble in aqueous buffers. For maximum solubility in aqueous buffers, erythromycin should first be dissolved in ethanol and then diluted with the aqueous buffer of choice. Erythromycin has a solubility of approximately 0.5 mg/ml in a 1:1 solution of ethanol:PBS (pH 7.2) using this method. We do not recommend storing the aqueous solution for more than one day.

Description

Erythromycin is a macrolide antibiotic that inhibits bacterial protein synthesis by targeting the 50S ribosomal subunit, blocking the progression of nascent polypeptide chains.¹ It is active against a host of bacterial genera, including Streptococcus, Staphylococcus, and Haemophilus (MIC₉₀s = 0.015-2.0 mg/l).² Erythromycin (10-40 mg/kg) dose-dependently inhibits the growth of S. aureus in a mouse model of thigh infection.³ It also inhibits the cytochrome P450 (CYP450) isoform CYP3A4 in vitro with IC₅₀ values of 33 and 27.3 μ M for α -hydroxytriazolam and 4-hydroxytriazolam formation, respectively, following administration of triazolam, which is known to be metabolized primarily by CYP3A4.^{4,5} Formulations containing erythromycin have been used in the treatment of bacterial respiratory and skin infections, pertussis, and a variety of other bacterial infections.

References

- 1. Wilson, D.N. Crit. Rev. Biochem. Mol. Biol. 44(6), 393-433 (2009).
- 2. Kanatani, M.S. and Guglielmo, B.J. Western J. Med. 160(1), 31-37 (1994).
- 3. Hoogeterp, J.J., Mattie, H., and van Furth, R. Scand. J. Infect. Dis. 25(1), 123-132 (1993).
- 4. Westphal, J.F. Br. J. Clin. Pharmacol. 50(4), 285-295 (2000).
- 5. Greenblatt, D.J., von Moltke, L.L., Harmatz, J.S., et al. Clin. Pharmacol. Ther. 64(3), 278-285 (1998).

WARNING THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFFTY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

uyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 05/22/2018

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897 [734] 971-3335 FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.CAYMANCHEM.COM