

Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com

PRODUCT INFORMATION



Norgestimate

Item No. 16547

	AS Registry No.: rmal Name:	35189-28-7 (17α)-17-(acetyloxy)-13-ethyl- 18,19-dinorpregn-4-en-20-yn-3-	°
Synonyms: MF: FW:		one 3-oxime Dexnorgestrel acetime, ORF 10131 C ₂₃ H ₃₁ NO ₃ 369.5	
Su	pplied as:	A crystalline solid	N V V
	orage: ability:	-20°C ≥2 years	
Information represents the preduct exections. Databased in a physical results are previded on each participate of analysis			

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

Norgestimate is supplied as a crystalline solid. A stock solution may be made by dissolving the norgestimate in the solvent of choice. Norgestimate is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide (DMF), which should be purged with an inert gas. The solubility of norgestimate in these solvents is approximately 2, 14, and 16 mg/ml, respectively.

Norgestimate is sparingly soluble in aqueous buffers. For maximum solubility in aqueous buffers, norgestimate should first be dissolved in DMF and then diluted with the aqueous buffer of choice. Norgestimate has a solubility of approximately 0.3 mg/ml in a 1:2 solution of DMF:PBS (pH 7.2) using this method. We do not recommend storing the aqueous solution for more than one day.

Description

Norgestimate is a progestin and a prodrug form of norelgestromin and levonorgestrel (Item No. 10006318).^{1,2} It binds to progestin receptors (mean IC_{50} = 3.48 nM for rabbit uterine receptors) and is selective for progestin over androgen receptors (mean IC_{50} = 764 nM for rat prostatic receptors) in radioligand binding assays.³ Norgestimate maintains pregnancy in ovariectomized female rats in a dose-dependent manner.¹ It inhibits ovulation in 100% of female rats when administered at a dose of 0.5 mg/kg. Norgestimate (2 mg/kg) completely inhibits estrone-induced vaginal cornification in ovariectomized female rats, indicating antiestrogenic activity. Formulations containing norgestimate in combination with ethinyl estradiol have been used as oral contraceptives and in the treatment of acne vulgaris.

References

- 1. Hahn, D.W., Allen, G.O., and McGuire, J.L. The pharmacological profile of norgestimate, a new orally active progestin. Contraception 16(5), 541-553 (1977).
- 2. Stanczyk, F.Z. All progestins are not created equal. Steroids 68(10-13), 879-890 (2003).
- 3. Phillips, A., Demarest, K., Hahn, D.W., et al. Progestational and androgenic receptor binding affinities and in vivo activities of norgestimate and other progestins. Contraception 41(4), 399-410 (1990).

WARNING THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFFTY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 09/24/2020

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897 [734] 971-3335 FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.CAYMANCHEM.COM