



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



Goat anti-Human (H+L) conjugated to HRP

Product Number HRP-110
Lot # HRP023-16-001
Amount 1 mg
Store at 2-8°C

Form/Shipping & Storage

Supplied as a lyophilized powder. Upon receipt, store at 2-8°C in the dark, do not freeze. Proteins are sensitive to freeze-thaw cycles.

Handling

Reconstitute to 1.0 ml with distilled deionized water vortex gently and allow vial to sit on ice for 20 minutes. We recommend that the investigator determine the appropriate working concentration for their specific application. Avoid exposure to heat and light.

Buffer

Upon rehydration with 1.0 ml distilled deionized water; the product is in 100 mM NaPO₄ (pH 7.4) + 100 mM NaCl + 1.5% BSA, 0.01% Thimerosal + 50mM Sucrose

Stability

Lyophilized material is stable for up to one year. After product has been reconstituted, product should be stored at 2-8°C in the dark and be used within 6 months. If further dilution of the conjugate is required, use diluted material within one week.

Note

For research use only, not for diagnostic or therapeutic use.

Concentration

After reconstitution - 1 mg/mL
13.58 μ M - HRP
3.16 μ M - Goat anti-Human

References:

Nygard NR, Bono C, Brown LR, Gorka J, Giacoletto KS, Schaiff WT, Graham MB, McCourt DW, Kabeer M, Braciale VL, et al. Antibody recognition of an immunogenic influenza hemagglutinin-human leukocyte antigen class II complex. *J Exp Med.* 1991 Jul 1;174(1):243-51.

Rogers BE, Chaudhuri TR, Reynolds PN, Della Manna D, Zinn KR. Non-invasive gamma camera imaging of gene transfer using an adenoviral vector encoding an epitope-tagged receptor as a reporter. *Gene Ther.* 2003 Jan;10(2):105-14.