

# Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



## Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

## Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



# **Product Information**



# 15(S)-15-methyl Prostaglandin F<sub>20</sub>

Item No. 16743

CAS Registry No: 35700-23-3

Formal Name:  $9\alpha,11\alpha,15S$ -trihydroxy-15-methyl-

prosta-5Z,13E-dien-1-oic acid

Synonyms: Carboprost, U-32921

MF:  $C_{21}H_{36}O_{5}$ FW: 368.5 **Purity:** ≥98%

Stability: ≥1 year at -20°C

Supplied as: A solution in methyl acetate

# COOH

#### **Laboratory Procedures**

15(S)-15-methyl Prostaglandin  $F_{2\alpha}$  (15(S)-15-methyl PGF $_{2\alpha}$ ) is a metabolically stable analog of PGF $_{2\alpha}$ . For long term storage, we suggest that 15(S)-15-methyl PGF $_{2a}$  be stored as supplied at -20°C. It will be stable for at least one year.

15(S)-15-methyl  $PGF_{2\alpha}$  is supplied as a solution in methyl acetate. To change the solvent, simply evaporate the methyl acetate under a gentle stream of nitrogen and immediately add the solvent of choice. Solvents such as ethanol, DMSO, and dimethyl formamide purged with an inert gas can be used. The solubility of 15(S)-15-methyl  $PGF_{2\alpha}$  in these solvents is approximately 100 mg/ml. 15(S)-15-methyl PGF $_{2\alpha}$  is stable for at least six months in these solvents if stored at -20°C. Further dilutions of the stock solution into aqueous buffers or isotonic saline should be made prior to performing biological experiments. Ensure that the residual amount of organic solvent is insignificant, since organic solvents may have physiological effects at low concentrations. If an organic solvent-free solution of 15(S)-15-methyl PGF<sub>2 $\alpha$ </sub> is needed, it can be prepared by evaporating the methyl acetate and directly dissolving the neat oil in aqueous buffers. The solubility of 15(S)-15-methyl PGF $_{2a}$  in PBS (pH 7.2) is approximately 10 mg/ml. We do not recommend storing the aqueous solution

15(S)-15-methyl  $PGF_{2\alpha}$  is a potent uterine stimulant and abortifacient which can be administered intramuscularly to induce labor. 1-3 It induces luteolysis and reduces serum progesterone concentrations when given as an intramuscular injection in non-pregnant rhesus monkeys.4

#### References

- 1. Carlan, S.J., Gushwa, J.P., O'Brien, W.F., et al. Effect of intramuscular 15-methyl prostaglandin F<sub>2α</sub> after secondtrimester delivery. Obstet. Gynecol. 89, 5-9 (1997).
- 2. Lindblom, B., Källfelt, B., Hahlin, M., et al. Local prostaglandin F<sub>2a</sub> injection for termination of ectopic pregnancy. Lancet, 776-777 (1987).
- 3. Adaikan, P.G., Prasad, R.N.V., Kottegoda, S.R., et al. Effect of indomethacin on the injection-abortion interval of 15(S)15 methyl PGF<sub>20</sub>-induced mid-trimester abortions - a randomized study. *Prostaglandins Leukot. Med.* **27**, 161-
- 4. Wilks, J.W. Inhibition of the monkey corpus luteum with 15-methyl prostaglandins. Prostaglandins 20, 793-805 (1980).

#### Related Products

For a list of related products please visit: www.caymanchem.com/catalog/16743

WARNING: This product is for laboratory research only: not for administration to humans. Not for human or veterinary DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until information to the contrary becomes available. Do not ingest, swallow, or inhale. Do not get in eyes, on skin, or on clothing. Wash thoroughly after handling. This information contains some, but not all, of the information required for the safe and proper use of this material. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

#### WARRANTY AND LIMITATION OF REMEDY

Cayman Chemical Company makes no warranty or guarantee of any kind, whether written or oral, expressed or implied, including without limitation, any warranty of fitness for a particular purpose, suitability and merchantability, which extends beyond the description of the chemicals hereof. Cayman warrants only to the original customer that the material will meet our specifications

the time of delivery.

Cayman will carry out its delivery obligations with due care and skill. Thus, in no event will Cayman have any obligation or liability, whether in tort (including negligence) or in contract, for direct, indirect, incidental or consequential damages, even if Cayman is informed about their possible existence.

This limitation of liability does not apply in the case of intentional acts or negligence of Cayman, its directors or its employees.

Buyer's exclusive remedy and Cayman's sole liability hereunder shall be limited to a refund of the purchase price, or at Cayman's option, the replacement, at no cost to Buyer, of all material that

Duyer's exclusive related and caylinais son analysis.

Solid refund or replacement is conditioned on Buyer giving written notice to Cayman within thirty (30) days after arrival of the material at its destination. Failure of Buyer to give said notice within thirty (30) days shall constitute a waiver by Buyer of all claims hereunder with respect to said material.

For further details, please refer to our Warranty and Limitation of Remedy located on our website and in our catalog.

Copyright Cayman Chemical Company, 06/18/2013

(734) 971-3640

(800) 364-9897

(734) 971-3335

custserv@caymanchem.com

www.caymanchem.com

Cayman Chemical

**Mailing address** 1180 E. Ellsworth Road Ann Arbor, MI 48108 USA