



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)


[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# PRODUCT INFORMATION



## 16-phenyl tetranor Prostaglandin F<sub>2α</sub> Item No. 16770

**CAS Registry No.:** 38315-48-9  
**Formal Name:** (5Z)-7-[(1R,2R,3R,5S)-3,5-dihydroxy-2-[[1E,3S)-3-hydroxy-4-phenyl-1-buten-1-yl]cyclopentyl]-5-heptenoic acid  
**Synonym:** 16-phenyl tetranor PGF<sub>2α</sub>  
**MF:** C<sub>22</sub>H<sub>30</sub>O<sub>5</sub>  
**FW:** 374.5  
**Purity:** ≥99%  
**Supplied as:** A crystalline solid  
**Storage:** -20°C  
**Stability:** ≥2 years



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

### Laboratory Procedures

16-phenyl tetranor Prostaglandin F<sub>2α</sub> (16-phenyl tetranor PGF<sub>2α</sub>) is supplied as a crystalline solid. A stock solution may be made by dissolving the 16-phenyl tetranor PGF<sub>2α</sub> in the solvent of choice, which should be purged with an inert gas. 16-phenyl tetranor PGF<sub>2α</sub> is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide. The solubility of 16-phenyl tetranor PGF<sub>2α</sub> in these solvents is approximately 100 mg/ml.

Further dilutions of the stock solution into aqueous buffers or isotonic saline should be made prior to performing biological experiments. Ensure that the residual amount of organic solvent is insignificant, since organic solvents may have physiological effects at low concentrations. Organic solvent-free aqueous solutions of 16-phenyl tetranor PGF<sub>2α</sub> can be prepared by directly dissolving the crystalline solid in aqueous buffers. The solubility of 16-phenyl tetranor PGF<sub>2α</sub> in PBS (pH 7.2) is approximately 0.8 mg/ml. We do not recommend storing the aqueous solution for more than one day.

### Description

16-phenyl tetranor PGF<sub>2α</sub> is a metabolically stable analog of PGF<sub>2α</sub>. The affinity of 16-phenyl tetranor PGF<sub>2α</sub> at the FP receptor of ovine luteal cells is poor (8.7%) compared to PGF<sub>2α</sub>.<sup>1</sup>

### Reference

1. Balapure, A.K., Rexroad, C.E., Jr., Kawada, K., *et al.* Structural requirements for prostaglandin analog interaction with the ovine corpus luteum prostaglandin F<sub>2α</sub> receptor. *Biochem. Pharmacol.* **38(14)**, 2375-2381 (1989).

#### WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

#### SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the [complete](#) Safety Data Sheet, which has been sent via email to your institution.

#### WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 05/03/2023

#### CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD  
ANN ARBOR, MI 48108 · USA

**PHONE:** [800] 364-9897  
[734] 971-3335

**FAX:** [734] 971-3640

CUSTSERV@CAYMANCHEM.COM  
WWW.CAYMANCHEM.COM