



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 


PRODUCT INFORMATION



20-ethyl Prostaglandin F_{2α}

Item No. 16940

CAS Registry No.: 36950-85-3
Formal Name: 9α,11α,15S-trihydroxy-20a,20b-dihomoprostanoic acid
Synonyms: ICI 74205, 20-ethyl PGF_{2α}
MF: C₂₂H₃₈O₅
FW: 382.5
Purity: ≥98%
Supplied as: A solution in methyl acetate
Storage: -20°C
Stability: ≥1 year



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

20-ethyl Prostaglandin F_{2α} (20-ethyl PGF_{2α}) is supplied as a solution in methyl acetate. To change the solvent, simply evaporate the methyl acetate under a gentle stream of nitrogen and immediately add the solvent of choice. Solvents such as ethanol, DMSO, and dimethyl formamide (DMF) purged with an inert gas can be used. The solubility of 20-ethyl PGF_{2α} in ethanol and DMF is approximately 33 mg/ml and approximately 14 mg/ml in DMSO.

Further dilutions of the stock solution into aqueous buffers or isotonic saline should be made prior to performing biological experiments. Ensure that the residual amount of organic solvent is insignificant, since organic solvents may have physiological effects at low concentrations. If an organic solvent-free solution of 20-ethyl PGF_{2α} is needed, it can be prepared by evaporating the methyl acetate and directly dissolving the neat oil in aqueous buffers. The solubility of 20-ethyl PGF_{2α} in PBS (pH 7.2) is approximately 3 mg/ml. We do not recommend storing the aqueous solution for more than one day.

Description

20-ethyl PGF_{2α} is an analog of PGF_{2α} in which the ω-chain has been extended by the addition of two more methylene carbon atoms. It is therefore a modified version of the clinically approved glaucoma medication unoprostone.¹ Unoprostone also contains lower side chain modifications (13,14-dihydro-15-keto) which severely limit its affinity for FP receptors, contributing to its lack of potency as a medication. 20-ethyl PGF_{2α} retains the natural 15(S) allylic hydroxyl in the lower side chain, which may improve its potency as an intraocular hypotensive agent compared to unoprostone. The 2 carbon extension in 20-ethyl-PGF_{2α} increases the K_i (120 nM) for the FP receptor from bovine corpus luteum only about 2.5-fold compared to PGF_{2α} (50 nM).² *In vivo* effects may be prolonged using 20-ethyl PGF_{2α}, as the activity of 15-hydroxy PGDH using 20-ethyl PGF_{2α} as a substrate is only 35% of the activity observed with PGF_{2α}.^{2,3}

References

1. Hara, M. and Spencer, C.M. Unoprostone (isopropyl unoprostone). *Drugs Aging* **9**(3), 213-218 (1996).
2. Powell, W.S., Hammarström, S., Samuelsson, B., *et al.* Interactions between prostaglandin analogues and a receptor in bovine *Corpora lutea*. Correlation of dissociation constants with luteolytic potencies in hamsters. *Eur. J. Biochem.* **59**(1), 271-276 (1975).
3. Sun, F.F., Armour, S.B., Bockstanz, V.R., *et al.* Studies on 15-hydroxyprostaglandin dehydrogenase from monkey lung. *Advances in prostaglandin and thromboxane research*. Samuelsson, B. and Paoletti, R., editors, Raven Press (1976).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 12/02/2021

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897
[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM