



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 


PRODUCT INFORMATION



(±)13(14)-DiHDPA

Item No. 18175

CAS Registry No.: 1345275-24-2
Formal Name: 13,14-dihydroxy-4Z,7Z,10Z,16Z,19Z-docosapentaenoic acid
Synonyms: 13(14)-DiHDPA, 13(14)-DiHDPE, 13,14-DiHDPE
MF: C₂₂H₃₄O₄
FW: 362.5
Purity: ≥98%
Supplied as: A solution in ethanol
Storage: -20°C
Stability: ≥1 year



NOTE: Relative stereochemistry shown in chemical structure

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

(±)13(14)-DiHDPA is supplied as a solution in ethanol. To change the solvent, simply evaporate the ethanol under a gentle stream of nitrogen and immediately add the solvent of choice. Solvents such as ethanol, DMSO, and dimethyl formamide purged with an inert gas can be used. The solubility of (±)13(14)-DiHDPA in these solvents is approximately 50 mg/ml.

Further dilutions of the stock solution into aqueous buffers or isotonic saline should be made prior to performing biological experiments. Ensure that the residual amount of organic solvent is insignificant, since organic solvents may have physiological effects at low concentrations. If an organic solvent-free solution of (±)13(14)-DiHDPA is needed, it can be prepared by evaporating the ethanol and directly dissolving the neat oil in aqueous buffers. The solubility of (±)13(14)-DiHDPA in PBS, pH 7.2, is approximately 0.25 mg/ml. We do not recommend storing the aqueous solution for more than one day.

Description

(±)13(14)-DiHDPA is a metabolite of docosahexaenoic acid (DHA; Item No. 90310) that is produced via oxidation by cytochrome P450 epoxygenases.¹ Epoxygenase metabolites of DHA, including (±)13(14)-DiHDPA, are reported to inhibit angiogenesis, tumor growth, and metastasis.²

References

1. VanRollins, M., Baker, R.C., Sprecher, H., *et al.* Oxidation of docosahexaenoic acid by rat liver microsomes. *J. Biol. Chem.* **259**, 5776-5783 (1984).
2. Zhang, G., Panigrahy, D., Mahakian, L.M., *et al.* Epoxy metabolites of docosahexaenoic acid (DHA) inhibit angiogenesis, tumor growth, and metastasis. *Proc. Natl. Acad. Sci. USA* **110(16)**, 6530-6535 (2013).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 05/16/2022

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM