

Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com

PRODUCT INFORMATION



Cardamonin

Item No. 18310

CAS Registry No.:	19309-14-9	
Formal Name:	(2E)-1-(2,4-dihydroxy-6-	
	methoxyphenyl)-3-phenyl-2- propen-1-one	ОН
Synonyms:	Alpinetin chalcone, Cardamomin	\downarrow \downarrow \land \land
MF:	C ₁₆ H ₁₄ O ₄	
FW:	270.3	
Purity:	≥98%	но
Stability:	≥2 years at -20°C	
Supplied as:	A crystalline solid	
UV/Vis.:	λ _{max} : 343 nm	

Laboratory Procedures

For long term storage, we suggest that cardamonin be stored as supplied at -20°C. It should be stable for at least two years.

Cardamonin is supplied as a crystalline solid. A stock solution may be made by dissolving the cardamonin in the solvent of choice. Cardamonin is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide (DMF), which should be purged with an inert gas. The solubility of cardamonin in ethanol is approximately 1 mg/ml and approximately 25 mg/ml in DMSO and DMF.

Cardamonin is sparingly soluble in aqueous buffers. For maximum solubility in aqueous buffers, cardamonin should first be dissolved in DMSO and then diluted with the aqueous buffer of choice. Cardamonin has a solubility of approximately 0.2 mg/ml in a 1:4 solution of DMSO:PBS (pH 7.2) using this method. We do not recommend storing the aqueous solution for more than one day.

Description

Cardamonin, isolated from the fruits of Alpinia species, is a chalconoid with anti-inflammatory and antitumor activity.^{1,2} It has been shown to suppress nitric oxide and prostaglandin E_2 synthesis, to suppress cyclooxygenase-2 expression, and to inhibit NF- κ B signaling.^{1,2} It can also target the Bcl-2 protein, inducing apoptosis in cancer cells.¹

References

- 1. Yadav, V.R., Prasad, S., Sung, B., et al. The role of chalcones in suppression of NF-κB-mediated inflammation and cancer. Int. Immunopharmacol. 11(3), 295-309 (2011).
- 2. Sung, B., Prasad, S., Yadav, V.R., et al. Cancer cell signaling pathways targeted by spice-derived nutraceuticals. Nutr. Cancer 64(2), 173-197 (2012).

WARNING THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFFTY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 11/05/2015

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897 [734] 971-3335 FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.CAYMANCHEM.COM