



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 


PRODUCT INFORMATION



1,6-Anhydro-D-galactose

Item No. 18895

CAS Registry No.: 644-76-8
Formal Name: 1,6-anhydro-β-D-galactopyranose
Synonyms: 1,6-Anhydro-β-D-galactopyranose, NSC 1376
MF: C₆H₁₀O₅
FW: 162.1
Purity: ≥95%
Stability: ≥2 years at -20°C
Supplied as: A crystalline solid



Laboratory Procedures

For long term storage, we suggest that 1,6-anhydro-D-galactose be stored as supplied at -20°C. It should be stable for at least two years.

1,6-Anhydro-D-galactose is supplied as a crystalline solid. A stock solution may be made by dissolving the 1,6-anhydro-D-galactose in the solvent of choice. 1,6-Anhydro-D-galactose is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide (DMF), which should be purged with an inert gas. The solubility of 1,6-anhydro-D-galactose in ethanol and DMSO is approximately 10 mg/ml and approximately 20 mg/ml in DMF.

Further dilutions of the stock solution into aqueous buffers or isotonic saline should be made prior to performing biological experiments. Ensure that the residual amount of organic solvent is insignificant, since organic solvents may have physiological effects at low concentrations. Organic solvent-free aqueous solutions of 1,6-anhydro-D-galactose can be prepared by directly dissolving the crystalline solid in aqueous buffers. The solubility of 1,6-anhydro-D-galactose in PBS, pH 7.2, is approximately 10 mg/ml. We do not recommend storing the aqueous solution for more than one day.

Description

1,6-Anhydro-D-galactose is a carbohydrate found in liquid smoke flavorings that is used as a tracer of the contribution of biomass burning to total atmospheric particulate matter.¹⁻³

References

1. Cordell, R.L., White, I.R., and Monks, P.S. Validation of an assay for the determination of levoglucosan and associated monosaccharide anhydrides for the quantification of wood smoke in atmospheric aerosol. *Anal. Bioanal. Chem.* **406**(22), 5283-5292 (2014).
2. Alañón, M.E., Rubio, H., Dkáz-Maroto, M.C., *et al.* Monosaccharide anhydrides, new markers of toasted oak wood used for aging wines and distillates. *Food Chem.* **119**, 505-512 (2010).
3. Guillén, M.D. and Manzanos, M.J. Study of the volatile composition of an aqueous oak smoke preparation. *Food Chem.* **79**, 283-292 (2002).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 02/23/2016

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897
[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM