

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



СООН

I-SAP

Item No. 19021

CAS Registry No.: 133538-58-6

Formal Name: (5Z)-7-[(1S,2S,3S,5R)-3-[[(4-iodophenyl)sulfonyl]

amino]-6,6-dimethylbicyclo[3.1.1]hept-2-yl]-5-

heptenoic acid

lodophenyl sulfonyl amino pinane Thromboxane A₂, Synonyms:

lodophenyl sulfonyl amino pinane TXA2

MF: $\mathrm{C}_{22}\mathrm{H}_{30}\mathrm{INO}_{4}\mathrm{S}$

FW: 531.4 **Purity:** ≥98%

UV/Vis.: λ_{max} : 249 nm

A solution in ethanol Supplied as:

-20°C Storage: Stability: ≥2 years

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.



I-SAP is supplied as a solution in ethanol. To change the solvent, simply evaporate the I-SAP under a gentle stream of nitrogen and immediately add the solvent of choice. Solvents such as DMSO and dimethyl formamide purged with an inert gas can be used. The solubility of I-SAP in these solvents is approximately 25 and 50 mg/ml, respectively.

Further dilutions of the stock solution into aqueous buffers or isotonic saline should be made prior to performing biological experiments. Ensure that the residual amount of organic solvent is insignificant, since organic solvents may have physiological effects at low concentrations. If an organic solvent-free solution of I-SAP is needed, it can be prepared by evaporating the ethanol and directly dissolving the neat oil in aqueous buffers. The solubility of I-SAP in PBS (pH 7.2) is approximately 100 µg/ml. We do not recommend storing the aqueous solution for more than one day.

Description

I-SAP is a high affinity TP receptor antagonist. At physiologic pH, I-SAP produces platelet shape change, but not aggregation, with an EC₅₀ value of 9.7 nM. I-SAP binds to human platelets with the maximum binding obtained between pH 6.5 and pH 7.4. In washed human platelets, the K_d for I-SAP is 468 pM at pH 7.4 and 490 pM at pH 6.5.1

Reference

1. Naka, M., Mais, D.E., Morinelli, T.A., et al. 7-[(1R,2S,3S,5R)-6,6-Dimethyl-3-(4-iodobenzenesulfonylamino) bicyclo[3.1.1]hept-2-yl]-5(Z)-heptenoic acid: A novel high-affinity radiolabeled antagonist for platelet thromboxane A₂/prostaglandin H₂ receptors. J. Pharmacol. Exp. Ther. **262(2)**, 632-637 (1992).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website

Copyright Cayman Chemical Company, 11/08/2021

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM