



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# PRODUCT INFORMATION



## 13,14-dihydro-15-keto Prostaglandin D<sub>2</sub>-d<sub>9</sub>

Item No. 19334


**Formal Name:** 9 $\alpha$ -hydroxy-11,15-dioxo-prost-5Z-en-1-oic-17,17,18,18,19,19,20,20,20-d<sub>9</sub> acid  
**Synonym:** 13,14-dihydro-15-keto-PGD<sub>2</sub>-d<sub>9</sub>  
**MF:** C<sub>20</sub>H<sub>23</sub>D<sub>9</sub>O<sub>5</sub>  
**FW:** 361.5  
**Chemical Purity:**  $\geq$ 98% (13,14-dihydro-15-keto PGD<sub>2</sub>)

**Deuterium Incorporation:**  $\geq$ 99% deuterated forms (d<sub>1</sub>-d<sub>9</sub>);  $\leq$ 1% d<sub>0</sub>

**Supplied as:** A solution in methyl acetate

**Storage:** -20°C

**Stability:**  $\geq$ 1 year



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

### Laboratory Procedures

13,14-dihydro-15-keto Prostaglandin<sub>2</sub>-d<sub>9</sub> (13,14-dihydro-15-keto PGD<sub>2</sub>-d<sub>9</sub>) is intended for use as an internal standard for the quantification of 13,14-dihydro-15-keto PGD<sub>2</sub>-d<sub>9</sub> (Item No. 12610) by GC- or LC-MS. The accuracy of the sample weight in this vial is between 5% over and 2% under the amount shown on the vial. If better precision is required, the deuterated standard should be quantitated against a more precisely weighed unlabeled standard by constructing a standard curve of peak intensity ratios (deuterated versus unlabeled).

13,14-dihydro-15-keto PGD<sub>2</sub>-d<sub>9</sub> is supplied as a solution in methyl acetate. To change the solvent, simply evaporate the methyl acetate under a gentle stream of nitrogen and immediately add the solvent of choice. Solvents such as ethanol, DMSO, and dimethyl formamide (DMF) purged with an inert gas can be used. The solubility of 13,14-dihydro-15-keto PGD<sub>2</sub>-d<sub>9</sub> in ethanol and DMF is approximately 50 mg/ml and 30 mg/ml in DMSO.

### Description

13,14-dihydro-15-keto PGD<sub>2</sub> is a metabolite of PGD<sub>2</sub> (Item No. 12010) which is formed through the 15-hydroxy PGDH pathway. 13,14-dihydro-15-keto PGD<sub>2</sub> was recently identified as a selective agonist for the CRTH<sub>2</sub>/DP<sub>2</sub> receptor.<sup>1</sup> It also inhibits ion flux in a canine colonic mucosa preparation.<sup>2</sup> In humans, 13,14-dihydro-15-keto PGD<sub>2</sub> is further metabolized to give 11 $\beta$ -hydroxy compounds which have also undergone  $\beta$ -oxidation of one or both side chains. Virtually no 13,14-dihydro-15-keto PGD<sub>2</sub> survives intact in the urine.<sup>3,4</sup>

### References

1. Rangachari, P.K. and Betti, P.A. Biological activity of metabolites of PGD<sub>2</sub> on canine proximal colon. *Am. J. Physiol.* **264**(5 Pt 1), G886-G894 (1993).
2. Hirai, H., Tanaka, K., Yoshie, O., et al. Prostaglandin D<sub>2</sub> selectivity induces chemotaxis in T helper type 2 cells, eosinophils, and basophils via seven-transmembrane receptor CRTH<sub>2</sub>. *J. Exp. Med.* **193**(2), 255-261 (2001).
3. Liston, T.E. and Roberts, L.J., II Metabolic fate of radiolabeled prostaglandin D<sub>2</sub> in a normal human male volunteer. *J. Biol. Chem.* **260**(24), 13172-13180 (1985).
4. Morrow, J.D., Prakash, C., Awad, J.A., et al. Quantification of the major urinary metabolite of prostaglandin D<sub>2</sub> by a stable isotope dilution mass spectrometric assay. *Anal. Biochem.* **193**(1), 142-148 (1991).

#### WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

#### SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

#### WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 05/11/2020

#### CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD  
ANN ARBOR, MI 48108 · USA

**PHONE:** [800] 364-9897

[734] 971-3335

**FAX:** [734] 971-3640

CUSTSERV@CAYMANCHEM.COM

WWW.CAYMANCHEM.COM