



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

PRODUCT INFORMATION



Cholera Toxin (azide free)

Item No. 19654

Overview and Properties

Synonyms:	Cholera toxin, CT
Contents:	Each vial contains 1 mg of cholera toxin (CT). When reconstituted to 0.5 ml with water, the buffer is 0.05 M Tris, 0.2 M NaCl, 0.001 M Na ₂ EDTA at pH 7.5
Storage:	2-8°C (as supplied)
Stability:	As supplied, 2 years from the QC date provided on the Certificate of Analysis, when stored properly
Purity:	≥95% (estimated by SDS-PAGE)
Protein	
Concentration:	Determined by absorbance at 280 nm using an extinction coefficient of 1.14 for a 1 mg/ml solution
Activity:	Exhibited hemagglutination activity at <2 ug/ml when assessed using G _{M1} fixed sheep red blood cells
Special conditions:	Handle gently; do not vortex; do not freeze

Description

Cholera toxin is produced by *V. cholerae* and is a multimeric enterotoxin that consists of a single A subunit surrounded by five B subunits.¹ Following binding to the membrane of enteric cells, the toxin is endocytosed and travels to the endoplasmic reticulum where the enzymatically active A subunit (CTA1) is translocated to the cytosol. Within the cytosol, CTA1 catalyzes the ADP-ribosylation of the G_sα protein, which activates adenylate cyclase, locking it in an 'on' position.² The increased intracellular concentration of cAMP causes hypersecretion of chloride and bicarbonate ions, which results in increased fluid secretion and severe diarrhea characteristic of the disease cholera.¹

References

1. Cho, J.A., Chinnapen, D.J.F., Amar, E., *et al.* Insights on the trafficking and retro-translocation of glycosphingolipid-binding bacterial toxins. *Front. Cell. Infect. Microbiol.* **2(51)** (2012).
2. van Heyningen, S. and King, C.A. Subunit A from cholera toxin is an activator of adenylate cyclase in pigeon erythrocytes. *Biochem. J.* **146(1)**, 269-271 (1975).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA
This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY
Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 08/10/2016

CAYMAN CHEMICAL
1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA
PHONE: [800] 364-9897
[734] 971-3335
FAX: [734] 971-3640
CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM