



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# PRODUCT INFORMATION



## LPS from *Escherichia coli* O55:B5

Item No. 19660

### Overview and Properties

---

<b>Contents:</b>	Each vial contains LPS isolated from <i>Escherichia coli</i> O55:B5, which was in water when lyophilized
<b>Storage:</b>	2-8°C (as supplied)
<b>Stability:</b>	≥2 years
<b>Special conditions:</b>	This product is pyrogenic. Avoid accidental autoinoculation by exercising extreme care when handling in conjunction with any injection device.

### Laboratory Procedures

---

LPS is dispersible in aqueous solvents at concentrations of 1 mg/ml. To achieve suspension in water, heating to about 50°C with intermittent vortexing or sonication is recommended. Allow ample time for dispersion to occur. The use of 0.5 % triethylamine aids in dispersion. Triethylamine is very basic and may be neutralized with Tris HCl to avoid hydrolysis of the LPS fatty acid chains.

### Description

---

LPS from *Escherichia coli* O55:B5 is a form of lipopolysaccharide (LPS) extracted from wild-type S-form *E. coli* serotype O55:B5. It is commonly used to activate toll-like receptor 4 (TLR4) on leukocytes, eliciting inflammatory signaling in isolated cells and *in vivo*.<sup>1-4</sup> This form of LPS includes three regions, the O-antigenic polysaccharide chain, the core oligosaccharide, and lipid A.

### References

---

1. Brauer, R.B., Gegenfurtner, C., Neumann, B., *et al.* Endotoxin-induced lung inflammation is independent of the complement membrane attack complex. *Infect. Immun.* **68(3)**, 1626-1632 (2000).
2. Jungi, T.W., Sager, H., Adler, H., *et al.* Serum factors, cell membrane CD14, and  $\beta$ 2 integrins are not required for activation of bovine macrophages by lipopolysaccharide. *Infect. Immun.* **65(9)**, 3577-3584 (1997).
3. Li, C.C., Munitic, I., Mittelstadt, P.R., *et al.* Suppression of dendritic cell-derived IL-12 by endogenous glucocorticoids is protective in LPS-induced sepsis. *PLoS Biol.* **13(10)**, (2015).
4. Migale, R., Herbert, B.R., Lee, Y.S., *et al.* Specific lipopolysaccharide serotypes induce differential maternal and neonatal inflammatory responses in a murine model of preterm labor. *Am. J. Pathol.* **185(9)**, 2390-2401 (2015).

WARNING  
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA  
This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY  
Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 05/15/2020

CAYMAN CHEMICAL  
1180 EAST ELLSWORTH RD  
ANN ARBOR, MI 48108 · USA  
PHONE: [800] 364-9897  
[734] 971-3335  
FAX: [734] 971-3640  
CUSTSERV@CAYMANCHEM.COM  
WWW.CAYMANCHEM.COM