

Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com

PRODUCT INFORMATION



3-hydroxy Palmitic Acid

Item No. 19934

CAS Registry No.:	2398-34-7
Formal Name:	3-hydroxy-hexadecanoic acid
Synonyms:	β-hydroxy Hexadecanoic Acid, OH
	β-hydroxy Palmitic Acid, \square
	DL-3-hydroxy Palmitic Acid HO \rightarrow \rightarrow \rightarrow
MF:	C ₁₆ H ₃₂ O ₃
FW:	272.4
Purity:	≥95%
Supplied as:	A crystalline solid
Storage:	-20°C
Stability:	As supplied, 2 years from the QC date provided on the Certificate of Analysis, when stored properly

Laboratory Procedures

3-hydroxy Palmitic acid is supplied as a crystalline solid. A stock solution may be made by dissolving the 3-hydroxy palmitic acid in the solvent of choice. 3-hydroxy Palmitic acid is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide (DMF), which should be purged with an inert gas. The solubility of 3-hydroxy palmitic acid in ethanol is approximately 2.5 mg/ml and approximately 20 mg/ml in DMSO and DMF.

3-hydroxy Palmitic acid is sparingly soluble in aqueous buffers. For maximum solubility in aqueous buffers, 3-hydroxy palmitic acid should first be dissolved in DMSO and then diluted with the aqueous buffer of choice. 3-hydroxy Palmitic acid has a solubility of approximately 0.33 mg/ml in a 1:2 solution of DMSO:PBS (pH 7.2) using this method. We do not recommend storing the aqueous solution for more than one day.

Description

3-hydroxy Palmitic acid is a form of the 16:0 lipid palmitic acid (Item No. 10006627). The lipid A part of lipopolysaccharides contain various 3-hydroxy fatty acids, making oxylipins such as 3-hydroxy palmitic acid useful as chemical markers of endotoxins.¹ In R. solanacearum, 3-hydroxy palmitic acid is converted by an S-adenosyl methionine-dependent methyltransferase to 3-hydroxy palmitic acid methyl ester, which acts as a quorum sensing signal molecule for post-transcriptional modulation of genes involved in virulence.² Long-chain 3-hydroxy fatty acids, such as 3-hydroxy palmitic acid, are also known to accumulate during long-chain 3-hydroxy-acyl-CoA dehydrogenase and mitochondrial trifunctional protein deficiencies.³ Such accumulation induces oxidative stress, leading to mitochondrial bioenergetics deregulation and eventual multi-organ dysfunction.³

References

- 1. Uhlig, S., Negård, M., Heldal, K. K., et al. J. Chromatogr. A. 1434, 119-236 (2016)
- 2. Shinohara, M., Nakajima, N., and Uehara, Y. J. Appl. Microbiol. 103(1), 152-162 (2007)
- 3. Tonin, A. M., Amaral, A. U., Busanello, E. N., et al. Biochim Biophys. Acta. 1842(9) 1658-1667 (2014)

WARNING THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFFTY DATA

al should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 08/08/2016

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897 [734] 971-3335 FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.CAYMANCHEM.COM