



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

PRODUCT INFORMATION



10-OAHSA

Item No. 19975

Formal Name: 10-(oleoyloxy)octadecanoic acid

MF: C₃₆H₆₈O₄


FW: 564.9

Purity: ≥95%

Supplied as: A solution in methyl acetate

Storage: -20°C

Stability: As supplied, 1 year from the QC date provided on the Certificate of Analysis, when stored properly



Laboratory Procedures

10-OAHSA is supplied as a solution in methyl acetate. To change the solvent, simply evaporate the methyl acetate under a gentle stream of nitrogen and immediately add the solvent of choice. Solvents such as ethanol, DMSO, and dimethyl formamide (DMF) purged with an inert gas can be used. The solubility of 10-OAHSA in ethanol and DMF is approximately 20 mg/ml and approximately 15 mg/ml in DMSO.

10-OAHSA is sparingly soluble in aqueous buffers. For maximum solubility in aqueous buffers, the ethanolic solution of 10-OAHSA should be diluted with the aqueous buffer of choice. 10-OAHSA has a solubility of approximately 0.5 mg/ml in a 1:1 solution of ethanol:PBS (pH 7.2) using this method. We do not recommend storing the aqueous solution for more than one day.

Description

10-OAHSA is a newly identified endogenous lipid that belongs to a collection of branched fatty acid esters of hydroxy fatty acids (FAHFAs). It is a FAHFA in which oleic acid (Item No. 90260) is esterified to 10-hydroxy stearic acid. Among the FAHFA family members, PAHSAs are the most abundant in the adipose tissue of glucose tolerant AG4OX mice, which overexpress the Glut4 glucose transporter specifically in adipose tissue.¹ As other FAHFAs improve glucose tolerance, stimulate insulin secretion, and have anti-inflammatory effects, 10-OAHSA may be a bioactive lipid with roles in metabolic syndrome and inflammation.¹

Reference

1. Yore, M.M., Syed, I., Moraes-Vieira, P.M., *et al.* Discovery of a class of endogenous mammalian lipids with anti-diabetic and anti-inflammatory effects. *Cell* **159**(2), 318-332 (2014).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 09/08/2016

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD

ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM

WWW.CAYMANCHEM.COM