

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



N-Oleoyl Valine

Item No. 20065

CAS Registry No.: 60374-41-6

N-[(9Z)-1-oxo-9-octadecen-1-yl]-L-valine Formal Name:

MF: $C_{23}H_{43}NO_{3}$ FW: 381.6 **Purity:** ≥98%

UV/Vis.: λ_{max} : 203 nm

Supplied as: A solution in methyl acetate

-20°C Storage: Stability: ≥2 years

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

N-Oleoyl valine is supplied as a solution in methyl acetate. To change the solvent, simply evaporate the methyl acetate under a gentle stream of nitrogen and immediately add the solvent of choice. Solvents such as ethanol, DMSO, and dimethyl formamide (DMF) purged with an inert gas can be used. The solubility of N-oleoyl valine in ethanol and DMSO is approximately 12 mg/ml and approximately 10 mg/ml in DMF.

N-Oleoyl valine is sparingly soluble in aqueous buffers. For maximum solubility in aqueous buffers, N-oleoyl valine should first be dissolved in ethanol and then diluted with the aqueous buffer of choice. N-Oleoyl valine has a solubility of approximately 0.2 mg/ml in a 1:4 solution of ethanol:PBS (pH 7.2) using this method. We do not recommend storing the aqueous solution for more than one day.

Description

N-Oleoyl valine is an endogenous N-acyl amine that acts as an antagonist at the transient receptor potential vanilloid type 3 (TRPV3) receptor, which is involved in thermoregulation.^{1,2} N-Oleoyl valine is increased in mice following cold exposure, up to at least 16 days. Acute lung injury in mice increases the concentration of N-oleoyl valine in lung tissue. N-acyl amines also promote mitochondrial uncoupling. 1

References

- 1. Raboune, S., Stuart, J.M., Leishman, E., et al. Novel endogenous N-acyl amides activate TRPV1-4 receptors, BV-2 microglia, and are regulated in brain in an acute model of inflammation. Front. Cell. Neurosci. 8(195), 1-11 (2014).
- 2. Broad, L.M., Mogg, A.J., Eberle, E., et al. TRPV3 in drug development. Pharmaceuticals 9(3), E55 (2016).
- 3. Long, J.Z., Svensson, K.J., Bateman, L.A., et al. The secreted enzyme PM20D1 regulates lipidated amino acid uncouplers of mitochondria. Cell 166, 1-12 (2016).
- 4. Balakrishna, S., Song, W., Achanta, S., et al. TRPV4 inhibition counteracts edema and inflammation and improves pulmonary function and oxygen saturation in chemically induced acute lung injury. Am. J. Physiol. Lung Cell. Mol. Physiol. 307(2), L158-L172 (2014).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website

Copyright Cayman Chemical Company, 05/19/2017

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM