



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 


PRODUCT INFORMATION



Phytosphingosine

Item No. 20217

CAS Registry No.: 554-62-1
Formal Name: 2S-amino-1,3S,4R-octadecanetriol
Synonym: D-ribo Phytosphingosine
MF: C₁₈H₃₉NO₃
FW: 317.5
Purity: ≥95%
Supplied as: A crystalline solid
Storage: -20°C
Stability: ≥2 years



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

Phytosphingosine is supplied as a crystalline solid. A stock solution may be made by dissolving the phytosphingosine in the solvent of choice. Phytosphingosine is soluble in organic solvents such as DMSO and dimethyl formamide, which should be purged with an inert gas. The solubility of phytosphingosine in these solvents is approximately 2 and 10 mg/ml, respectively. Phytosphingosine is also miscible in ethanol.

Description

Phytosphingosine is a sphingolipid with a hydroxyl group at the C4 position that is found mainly in fungi and plants but also in animals, including humans.^{1,2} It is metabolized to odd-numbered fatty acids with 2-hydroxy palmitic acid (Item No. 22679) as an intermediate.³ Phytosphingosine dose-dependently induces cell death of CHO cells and inhibits carbachol-induced activation of phospholipase D (PLD) in CHO cells transfected with *C. elegans* muscarinic acetylcholine receptors.⁴ It is essential in the heat stress response in *S. cerevisiae*.⁵

References

1. Dickson, R.C. Sphingolipid functions in *Saccharomyces cerevisiae*: Comparison to mammals. *Annu. Rev. Biochem.* **67**, 27-48 (1998).
2. Schürer, N.Y., Plewig, G., Elias, P.M., *et al.* Stratum corneum lipid function. *Dermatologica* **183(2)**, 77-94 (1991).
3. Kondo, N., Ohno, Y., Yamagata, M., *et al.* Identification of the phytosphingosine metabolic pathway leading to odd-numbered fatty acids. *Nat. Commun.* **5**, 5338 (2014)
4. Lee, J.S., Min, D.S., Park, C., *et al.* Phytosphingosine and C2-phytoceramide induce cell death and inhibit carbachol-stimulated phospholipase D activation in Chinese hamster ovary cells expressing the *Caenorhabditis elegans* muscarinic acetylcholine receptor. *FEBS. Lett.* **499(1-2)**, 82-86 (2001).
5. Hannun, Y.A., Luberto, C., and Argraves, K.M. Enzymes of sphingolipid metabolism: From modular to integrative signaling. *Biochemistry* **40(16)**, 4893-4903 (2001).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the [complete](#) Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 11/14/2017

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897
[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM