



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 


PRODUCT INFORMATION



Deoxycholic Acid

Item No. 20756

CAS Registry No.: 83-44-3
Formal Name: (3 α ,5 β ,12 α)-3,12-dihydroxy-cholan-24-oic acid
Synonyms: Cholanoic Acid, DCA, NSC 8797
MF: C₂₄H₄₀O₄
FW: 392.6
Purity: \geq 95%
Supplied as: A crystalline solid
Storage: -20°C
Stability: \geq 2 years



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

Deoxycholic acid (DCA) is supplied as a crystalline solid. A stock solution may be made by dissolving the DCA in the solvent of choice, which should be purged with an inert gas. DCA is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide (DMF). The solubility of DCA in ethanol and DMSO is approximately 20 mg/ml and approximately 30 mg/ml in DMF.

DCA is sparingly soluble in aqueous buffers. For maximum solubility in aqueous buffers, DCA should first be dissolved in DMF and then diluted with the aqueous buffer of choice. DCA has a solubility of approximately 0.5 mg/ml in a 1:1 solution of DMF:PBS (pH 7.2) using this method. We do not recommend storing the aqueous solution for more than one day.

Description

DCA is a secondary bile acid that is formed *via* microbial transformation of cholic acid (Item No. 20250) in the colon.¹ It can be conjugated to glycine or taurine (Item No. 27031) to produce glycodeoxycholic acid (GDCA; Item No. 20274) or taurodeoxycholic acid (TDCA; Item No. 15935), respectively, in hepatocytes.¹⁻³ DCA (0.2% v/v) inhibits spore germination induced by taurocholic acid (TCA; Item No. 16215) in seven *C. difficile* strains, as well as inhibits growth and decreases the cytotoxicity of *C. difficile* culture supernatants to Vero cells when used at a concentration of 0.02% v/v.¹ It inhibits ionizing radiation-induced p53-dependent transcription in a reporter assay using HCT116 cells when used at a concentration of 200 μ M.⁴ Fecal and intestinal tissue levels of DCA are increased in a rat model of high-fat diet-induced obesity compared with rats fed a normal diet.⁵ Increased serum DCA levels have been found in patients with colorectal cancer.⁶

References

1. Thanissery, R., Winston, J.A., and Theriot, C.M. Inhibition of spore germination, growth, and toxin activity of clinically relevant *C. difficile* strains by gut microbiota derived secondary bile acids. *Anaerobe* **45**, 86-100 (2017).
2. Schmid, A., Neumann, H., Karrasch, T., *et al.* Bile acid metabolome after an oral lipid tolerance test by liquid chromatography-tandem mass spectrometry (LC-MS/MS). *PLoS One* **11**(2), e0148869 (2016).
3. Šarenac, T.M. and Mikov, M. Bile acid synthesis: From nature to the chemical modification and synthesis and their applications as drugs and nutrients. *Front. Pharmacol.* **9**, 939 (2018).
4. Qiao, D., Gaitonde, S.V., Qi, W., *et al.* Deoxycholic acid suppresses p53 by stimulating proteasome-mediated p53 protein degradation. *Carcinogenesis* **22**(6), 957-964 (2001).
5. Lin, H., An, Y., Tang, H., *et al.* Alterations of bile acids and gut microbiota in obesity induced by high fat diet in rat model. *J. Agric. Food Chem.* **67**(13), 3624-3632 (2019).
6. Bayerdörffer, E., Mannes, G.A., Richter, W.O., *et al.* Increased serum deoxycholic acid levels in men with colorectal adenomas. *Gastroenterology* **104**(1), 145-151 (1993).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the [complete](#) Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 08/05/2020

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897
[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM