

# Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



# Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

# Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

## SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



# PRODUCT INFORMATION



OH

Н

Н

## 17α-Estradiol

Item No. 20776

CAS Registry No.: 57-91-0

(17α)-estra-1,3,5(10)-triene-3,17-diol Formal Name: Synonyms: Alfatradiol, α-Estradiol, 17-epi Estradiol,

NSC 20293, 17α-Oestradiol

MF:  $C_{18}H_{24}O_{2}$ 272.4 FW: ≥98% **Purity:** 

UV/Vis.:  $\lambda_{\text{max}}$ : 202, 281 nm Supplied as: A crystalline solid

Storage:

As supplied, 2 years from the QC date provided on the Certificate of Analysis, when Stability:

stored properly

#### **Laboratory Procedures**

 $17\alpha$ -Estradiol is supplied as a crystalline solid. A stock solution may be made by dissolving the  $17\alpha$ -estradiol in the solvent of choice.  $17\alpha$ -Estradiol is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide (DMF), which should be purged with an inert gas. The solubility of  $17\alpha$ -estradiol in ethanol is approximately 2.5 mg/ml and approximately 20 mg/ml in DMSO and DMF.

17α-Estradiol is sparingly soluble in aqueous buffers. For maximum solubility in aqueous buffers, 17α-estradiol should first be dissolved in DMSO and then diluted with the aqueous buffer of choice. 17α-Estradiol has a solubility of approximately 0.2 mg/ml in a 1:4 solution of DMSO:PBS (pH 7.2) using this method. We do not recommend storing the aqueous solution for more than one day.

#### Description

17α-Estradiol is the natural optical isomer of 17β-estradiol (Item No. 10006315), the major estrogen secreted by the premenopausal ovary.  $17\alpha$ -Estradiol is a less active isomer than its counterpart, displaying 100-fold lower estrogenic activity relative to  $17\beta$ -estradiol. It can inhibit  $5\alpha$ -reductase, which plays a role in regulating hair growth. 17a -estradiol is reported to activate the MAPK/ERK and PI3K-Akt signaling pathways via activation of the estrogen receptor-X and has been shown to be neuroprotective after an ischemic stroke and oxidative stress and in transgenic mice with Alzheimer's disease. 1,2

#### References

- 1. Toran-Allerand, C.D., Tinnikov, A.A., Singh, R.J., et al. 17α-estradiol: A brain-active estrogen? Endocrinology 146(9), 3843-3850 (2005).
- 2. Green, P.S., Bishop, J., and Simpkins, J.W. 17α-estradiol exerts neuroprotective effects on SK-N-SH cells. J. Neurosci. 17(2), 511-515 (1997).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

### WARRANTY AND LIMITATION OF REMEDY

subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website

Copyright Cayman Chemical Company, 12/21/2016

#### **CAYMAN CHEMICAL**

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM